arXiv:1203.1943v2 [math.RT] 1 Jul 2012

Monoid algebras of projection functors

Anna-Louise Grensing
Fakultät für Mathematik
Universität Bielefeld
D - 36615 Bielefeld
agrensin@math.uni-bielefeld.de

25th February 2013

Abstract

We study the monoid of so called projection functors P_S attached to simple modules S of a finite dimensional algebra, which appear naturally in the study of torsion pairs. We determine defining relations in special cases of path algebras. For the linearly oriented Dynkin quiver of Type A, we get an isomorphism to the monoid of non-decreasing parking functions. Moreover we give an explicit isomorphism between the monoid algebra of non-decreasing parking functions and a certain incidence algebra independent of the field.

1 Introduction

The study of Hom-orthogonal subcategories is a classical tool in the representation theory of finite dimensional algebras. The process of passing from the module category to orthogonal subcategories is essential as pointed out for example in [GL91], [Sch91]. Here the so called projection functors appear naturally. We start by defining them for arbitrary modules U over an associative finite dimensional unital algebra A over a field k. We denote by A-mod the category of finite dimensional left A-modules and by gen-U its full subcategory which consists of those modules isomorphic to a quotient of some $U^{\oplus d}$. We first define the endofunctor t_U on A-mod by sending a module M to its greatest submodule $\mathrm{t}_U(M)$ which lies in gen-U. This gives a subfunctor of the identity functor $\mathrm{id}_{A\text{-mod}}$, i.e. the embeddings ι_M of $\mathrm{t}_U(M)$ into M for every module M yield a natural transformation $\iota\colon \mathrm{t}_U \longrightarrow \mathrm{id}_{A\text{-mod}}$. We obtain the so called projection functor $\mathsf{P}_U\colon A\text{-mod} \longrightarrow A\text{-mod}$ by passing to the cokernel $(\pi_M)_{M\in A\text{-mod}}\colon \mathrm{id}_{A\text{-mod}} \longrightarrow \mathsf{P}_U$ of ι.
The question of describing the relations between such functors arises naturally. It fits into the general categorification programmme of realizing Lie-theoretic objects as functors on module categories [Maz10].
We concretrize this and concentrate just on the multiplicative interplay between certain projection functors. To create the framework we consider the monoid π_A generated by $\{\mathsf{P}_S \mid S\,\text{simple}\}$

up to natural isomorphism. For this we recall that the composition of endofunctors on A-mod induces a multiplication on the isomorphism classes of endofunctors. Hence π_A is the set of the isomorphism classes $[\mathsf{P}_{S_1}\mathsf{P}_{S_2}\dots\mathsf{P}_{S_r}]$ with $r \geq 0$ and $S_1, S_2, \dots, S_r$ simple A-modules together with that multiplication. In this sense we speak of monomials over $\{\mathsf{P}_S \mid S \,\text{simple}\,\}$, omit the brackets indicating the isomorphism classes and write 1 for the isomorphism class of the identity functor. To study this monoid and its monoid algebra a good way to start is finding a set of defining relations. We give some relations:

Proposition 1.1. *Let S and T be simple A-modules without any non-trivial self-extensions. Then the following relations hold:*

(a) $\mathsf{P}_S \circ \mathsf{P}_S = \mathsf{P}_S$

(b) If $\operatorname{Ext}^1_A(T, S) = 0$ *holds, we have* $\mathsf{P}_S \circ \mathsf{P}_T \circ \mathsf{P}_S = \mathsf{P}_T \circ \mathsf{P}_S \circ \mathsf{P}_T = \mathsf{P}_S \circ \mathsf{P}_T$.

As a consequence of the second relations we get $\mathsf{P}_S \circ \mathsf{P}_T = \mathsf{P}_T \circ \mathsf{P}_S$ if $\operatorname{Ext}^1_A(T, S) = 0 = \operatorname{Ext}^1_A(S, T)$ holds. Thus certain generators of π_A satisfy the braid relations of type A.
Now if A is the path algebra kQ of a finite acyclic quiver $Q = (Q_0, Q_1)$ over k, then each simple kQ-module S_t (attached to the vertex t) has no non-trivial self-extensions and hence the proposition holds for all generators of $\pi_Q := \pi_{kQ}$. We conjecture that these relations are defining for π_Q. To determine whether the above relations are defining ones we compare $k\pi_Q$ with the algebra B_Q, which we define by generators X_t with $t \in Q_0$ and the following relations:

- $X_s^2 = X_s$ for all $s \in Q_0$
- $X_s X_t X_s = X_t X_s X_t$ for all $s, t \in Q_0$
- $X_t X_s X_t = X_s X_t$ for all $s, t \in Q_0$, such that there is no arrow from t to s

There is an epimorphism ψ_Q of algebras from $\mathcal{B}_Q$ onto $k\pi_Q$ with $X_t \mapsto \mathsf{P}_{S_t} =: \mathsf{P}_t$ for all $t \in Q_0$ by the above relations of the projection functors since the dimension of $\operatorname{Ext}^1_{kQ}(S_t, S_s)$ coincides with the number of arrows from t to s. Note that if this epimorphism ψ_Q is an isomorphism, then π_Q is isomorphic to the Hecke-Kiselman semigroup associated with Q introduced in [GM11]. We introduce a method to detect when ψ is an isomorphism in section 3. So far we applied it successfully to tree quivers with a specific orientation including bipartite tree quivers, m-subspace quivers, star quivers and each Dynkin quiver of type A as well as to a couple of families of symmetrically shaped quivers. In this article we just discuss the former ones (see Prop. 3.7 and Theo. 3.9). As the relations are independent of the number of arrows unless there are none, one expects that multiple arrows have no impact. Moreover the relations are local as they just take direct neighbourhoods into account. We treat these aspects in section 3 for $\mathcal{B}_Q$ where it is obvious and for $k\pi_Q$.
In section 4 we study the algebra $\mathcal{B}_Q$ and compute its Gabriel quiver. By definition its underlying monoid is the Hecke-Kiselman semigroup associated with Q. These algebras emerge as finite dimensional (see Cor. 3.3) and basic (see Prop. 4.1) regardless of the representation type of Q. The Gabriel quiver can be described using the combinatorics given by the shape of the original quiver.

Theorem 1.2. *The simple modules E_M of $\mathcal{B}_Q$, hence the vertices of the Gabriel quiver $\Gamma(\mathcal{B}_Q)$, are parametrized by the subsets M of Q_0. Moreover there is at most one arrow between two vertices. More precisely we have $[\mathrm{E}_M] \rightarrow [\mathrm{E}_N] \in \Gamma(\mathcal{B}_Q)$ for two subsets M, N of Q_0 if and only if $M \setminus N$ and $N \setminus M$ are non-empty and if for each pair $(m, n) \in M \setminus N \times N \setminus M$ there is an arrow $m \rightarrow n$ in Q.*

Since the Hecke-Kiselman semigroup associated with an acyclic finite quiver is $\mathcal{J}$-trivial, these results can be deduced from [DHST11]. The author thanks Anne Schilling for pointing this reference out. Let Q_n denote the following linearly oriented Dynkin quiver of type A:

$$1 \longrightarrow 2 \longrightarrow \ \ldots \ \longrightarrow n$$

This family of quivers has a special role, we show in subsection 4.3:

Proposition 1.3. *Let Q be connected, finite and acyclic.*

(a) If $Q = Q_n$ then the Gabriel quiver of $\mathcal{B}_{Q_n}$ has exactly $n+1$ connected components.

(b) If Q is distinct from the linearly oriented Dynkin quiver of type A, then the Gabriel quiver of $\mathcal{B}_Q$ has exactly 3 connected components.

Therefore we devote section 5 to the algebra $\mathcal{B}_{Q_n}$, which is isomorphic to $k\pi_{Q_n}$ by section 3. It can be read almost independent. Now the monoid algebra of non-decreasing parking functions NDPF_{n+1} has the same defining relations as $\mathcal{B}_{Q_n}$ (see [HT09] or [GM11]) and is thus isomorphic to $\mathcal{B}_{Q_n}$. As shown in [HT09] it is isomorphic to the incidence algebra $\mathrm{Inc}\,(\mathcal{P}_n)$ of the product order $\geq_n$ on the powerset of $\underline{n\rfloor} := \{1, 2, \ldots, n\}$ if the underlying field k is the field of complex numbers $\mathbb{C}$. We recall, that for two subsets $K = \{k_1 < \ldots < k_r\}$ and $J = \{j_1 < \ldots < j_m\}$ of $\underline{n\rfloor}$ we have $K \geq_n J$ iff $m = r$ and $k_i \geq j_i$ holds for all $i \in \underline{r\rfloor}$. They use the representation theory of the symmetric group. In [DHST11] this is generalized using the $\mathcal{J}$-triviality of the underlying monoid. Here (see Main Theorem 5.3) we inductively construct an isomorphism from $\mathrm{Inc}\,(\mathcal{P}_n)$ to $\mathcal{B}_{Q_n}$ independent of the field – in fact it holds for a commutative ring – using the structure of the tower of algebras $\mathcal{B}_{Q_1} \subset \mathcal{B}_{Q_2} \subset \ldots$ and the action of π_{Q_n} on the (injective indecomposable) kQ_n-modules. We expect that a notion of non-decreasing parking functions for arbitrary quivers determines defining relations for π_Q.

Acknowledgments:
This work is part of my PhD-thesis [Paa11] supervised by Markus Reineke whom I thank for posing this question.

2 Relations

For every homomorphism $\varphi\colon M \longrightarrow N$ in A-mod we get the following commutative diagram defining $\mathsf{P}_U\varphi$, which we will call the diagram of φ induced by P_U, with exact rows, which we will call the short exact sequence of M resp. N induced by P_U:

$$\begin{tikzcd}
0 \arrow[r] & \mathrm{t}_U M \arrow[r, "\iota_M"] \arrow[d, "\mathrm{t}_U(\varphi) = \varphi|_{\mathrm{t}_U M}"'] & M \arrow[r, "\pi_M"] \arrow[d, "\varphi"'] & \mathsf{P}_U M = M/\mathrm{t}_U M \arrow[r] \arrow[d, dotted, "\mathsf{P}_U(\varphi)"'] & 0 \\
0 \arrow[r] & \mathrm{t}_U N \arrow[r, "\iota_N"] & N \arrow[r, "\pi_N"] & \mathsf{P}_U N = N/\mathrm{t}_U N \arrow[r] & 0
\end{tikzcd}$$

Other descriptions of the endofunctor t_U are useful. Obviously for every A-module M, the module $\mathrm{t}_U M$ is the sum over all submodules X of M lying in gen-U. Therefore $\mathrm{t}_U M$ is the image of the evaluation map $\mathrm{ev}_{U,M}\colon U \otimes_{\mathrm{End}_A(U)} \mathrm{Hom}_A(U, M) \longrightarrow M$ with $u \otimes \varphi \mapsto \varphi(u)$. Now if $U = S$ is simple then the module $\mathrm{t}_S M$ is isomorphic to some $S^{\oplus d}$ and $\mathrm{ev}\colon S \otimes_{\mathrm{End}_A(S)} \mathrm{Hom}_A(S, \_) \longrightarrow \mathrm{t}_S$ is thus even a natural isomorphism.

Proof of Proposition 1.1(a). If S has no non-trivial self-extensions, then t_S is a torsion radical, i.e. a subfunctor of the identity functor such that $\mathrm{t}_S(M/\mathrm{t}_S M) = 0$ holds for all $M \in A$-mod, and $\mathrm{im}\,\mathsf{P}_S = \ker \mathrm{Hom}_A(S, \_)$ holds. ☐

We will prove (b) of Proposition 1.1 with (and after) the following lemma.

Lemma 2.1. *Let S and T be two non-isomorphic simple modules. Then $\mathsf{P}_T \circ \mathsf{P}_S$ and $\mathsf{P}_{S\oplus T}$ are naturally isomorphic if and only if $\mathrm{Ext}^1_A(T, S) = 0$ holds.*

Proof. If there exists a non-split short exact sequence $0 \longrightarrow S \xrightarrow{f} X \longrightarrow T \longrightarrow 0$, then the functors $\mathsf{P}_T \circ \mathsf{P}_S$ and $\mathsf{P}_{S\oplus T}$ are not naturally isomorphic since we have:

$$\mathsf{P}_T \circ \mathsf{P}_S(X) = \mathsf{P}_T(X/f(S)) = 0 \neq T \cong X/f(S) = \mathsf{P}_{S\oplus T}(X)$$

Now we assume $\mathrm{Ext}^1_A(T, S) = 0$. Let M be a module. Since $\mathrm{t}_{S\oplus T} M$ is semi-simple it coincides with $\mathrm{t}_S M \oplus \mathrm{t}_T M$. For that reason the restriction of the canonical projection $\pi\colon M \longrightarrow M/\mathrm{t}_S M$ to $\mathrm{t}_{S\oplus T} M$ factors through $\pi'\colon \mathrm{t}_{S\oplus T} M \longrightarrow \mathrm{t}_T(\mathsf{P}_S M)$. By passing to the cokernels we get the following commutative diagram whose exact rows are induced by $\mathsf{P}_{S\oplus T}$ and P_T:

$$\begin{tikzcd}
0 \arrow[r] & \mathrm{t}_{S\oplus T} M \arrow[r] \arrow[d, "\pi'"] & M \arrow[r] \arrow[d, "\pi"] & \mathsf{P}_{S\oplus T} M = M/\mathrm{t}_{S\oplus T} M \arrow[r] \arrow[d, dotted, "\hat{\pi} = \hat{\pi}_M"] & 0 \\
0 \arrow[r] & \mathrm{t}_T(\mathsf{P}_S M) \arrow[r] & \mathsf{P}_S M \arrow[r] & \mathsf{P}_T \mathsf{P}_S M = (\mathsf{P}_S M)/(\mathrm{t}_T(M/\mathrm{t}_S M)) \arrow[r] & 0
\end{tikzcd}$$

To see that the epimorphism $\hat{\pi}_M$ is an isomorphism, it suffices to show that π' is an epimorphism because of $\ker \pi' \cong \ker \pi$. Since $\mathrm{Hom}_A(T, M)$ and $\mathrm{Hom}_A(T, \mathsf{P}_S M)$ are isomorphic by the assumptions, this follows from:

$$\mathrm{t}_T(\mathsf{P}_S M) \cong T \otimes_{\mathrm{End}(T)} \mathrm{Hom}_A(T, \mathsf{P}_S M) \cong T \otimes_{\mathrm{End}(T)} \mathrm{Hom}_A(T, M) \cong \mathrm{t}_T M \cong \pi'(\mathrm{t}_{S\oplus T} M)$$

Moreover we get for every morphism $\varphi\colon M \longrightarrow N$ the following cube,in which the sides on the left commute and in which thus the right square commutes. Thus $\hat{\pi}$ is a natural isomorphism.

$$\begin{tikzcd}
 & N \arrow[rr] \arrow[dd, dotted] & & \mathsf{P}_{S\oplus T}N \arrow[dd, "\hat{\pi}_N"] & \\
M \arrow[ur, "\varphi"] \arrow[rr] \arrow[dd] & & \mathsf{P}_{S\oplus T}M \arrow[ur, "\mathsf{P}_{S\oplus T}\varphi"] \arrow[rr] \arrow[dd, "\hat{\pi}_M"] & & 0 \\
 & \mathsf{P}_S N \arrow[rr, dotted] & & \mathsf{P}_T\mathsf{P}_S N & \\
\mathsf{P}_S M \arrow[ur, dotted, "\mathsf{P}_S\varphi"] \arrow[rr] & & \mathsf{P}_T\mathsf{P}_S M \arrow[ur, "\mathsf{P}_T\mathsf{P}_S\varphi"] & &
\end{tikzcd}$$

□

Proof of Proposition 1.1(b). By the previous lemma it suffices to show
(A) $\mathsf{P}_{S\oplus T}\circ\mathsf{P}_T \sim \mathsf{P}_S\circ\mathsf{P}_T$ and (B) $\mathsf{P}_S\circ\mathsf{P}_{S\oplus T} \sim \mathsf{P}_S\circ\mathsf{P}_T$.
Proof of (A). Since T has no non-trivial self-extensions and is as simple as S, a natural isomorphism is induced by:

$$\mathsf{t}_{S\oplus T}(\mathsf{P}_T M) \cong \mathsf{t}_S(\mathsf{P}_T M) \oplus \mathsf{t}_T(\mathsf{P}_T M) \cong \mathsf{t}_S(\mathsf{P}_T M)$$

Proof of (B). Let $\alpha\colon \mathsf{P}_T \longrightarrow \mathsf{P}_{S\oplus T}$ be the natural transformation given by the following composition α_M of the canonical epimorphism and isomorphism for every module M:

$$\begin{tikzcd}
\mathsf{P}_T M = M/\mathsf{t}_T M \arrow[r, two heads] \arrow[rr, bend left, "\alpha_M"] & (M/\mathsf{t}_T M)\Big/\Big((\mathsf{t}_S M \oplus \mathsf{t}_T M)/\mathsf{t}_T M\Big) \arrow[r, "\cong"] & \mathsf{P}_{S\oplus T}M
\end{tikzcd}$$

We claim, that the natural transformation $(\mathsf{P}_S\alpha_M)_{M\in A\text{-mod}}\colon \mathsf{P}_S\circ\mathsf{P}_T \longrightarrow \mathsf{P}_S\circ\mathsf{P}_{S\oplus T}$ is a natural isomorphism. For this we look at the diagram of α_M induced by P_S and consider its exact sequence of kernels and cokernels given by the snake lemma:

$$0 \to \ker \mathsf{t}_S\alpha_M \xrightarrow{\iota} \ker\alpha_M \longrightarrow \ker \mathsf{P}_S\alpha_M \longrightarrow \operatorname{coker} \mathsf{t}_S\alpha_M \longrightarrow 0 \longrightarrow \operatorname{coker}\mathsf{P}_S\alpha_M \to 0$$

Now $\ker\alpha_M \cong \mathsf{t}_S M \in \text{gen-}S$ is a submodule of $\mathsf{t}_S\mathsf{P}_T M$. Hence the monomorphism ι is an isomorphism. Moreover $\mathsf{t}_S\alpha_M$ is surjective. This is seen by using the assumptions on S, which yield the surjectivity of $\mathrm{Hom}_A(S,\alpha_M)$, and the natural isomorphism of t_S and $S\otimes\mathrm{Hom}_A(S,\_)$. □

3 The monoid algebras $k\pi_Q$ and $\mathcal{B}_Q$ for path algebras kQ

We now turn towards the class of finite dimensional path algebras. So let $Q = (Q_0, Q_1)$ be a finite acyclic quiver, i.e. Q is an oriented graph without oriented cycles and with finite sets Q_0 and Q_1

of vertices and arrows respectively. We will denote an arrow in Q by $\alpha\colon s \to t \in Q_1$ or $s \xrightarrow{\alpha} t$, and a path $t_n \xleftarrow{\beta_n} \dots \xleftarrow{\beta_2} t_1 \xleftarrow{\beta_1} t$ by $\beta_n \dots \beta_2\beta_1$. The path algebra of Q over a field k is denoted by kQ. The category kQ-mod and the category $Q\text{-}_k\mathrm{rep}$ of finite dimensional Q-representations over k are equivalent and we will not distinguish between them. Let $(Q_0^*, \boldsymbol{.})$ be the free monoid over Q_0, i.e. the monoid of words over the alphabet Q_0 with the concatenation $\boldsymbol{.}$ as multiplication. There is the canonical epimorphism $\rho\colon kQ_0^* \longrightarrow \mathcal{B}_Q$ with $q \mapsto X_q$ for all $q \in Q_0$. We will denote the image of a word $w \in Q_0^*$ under ρ by X_w. By P_w we denote the image of X_w under the canonical epimorphism $\psi_Q\colon \mathcal{B}_Q \longrightarrow k\pi_Q$.

Definition 3.1. *We call a subset W of Q_0^* an admissible normal form associated with Q if the following conditions hold:*

(1) $\{\emptyset\} \cup Q_0 \subseteq W$

(2) $B_W := \{X_w \mid w \in W\}$ *is closed under (right-) multiplication with the generators X_t of $\mathcal{B}_Q$.*

(3) For all words $v \neq w$ in W, there is a Q-representation V with $\mathsf{P}_v V \ncong \mathsf{P}_w V$.

Obviously the set $\{X_w \mid w \in Q_0^*\}$ is a k-linear generating system of $\mathcal{B}_Q$. This definition extracts suitable conditions on a subset of $\{X_v \mid v \in Q_0^*\}$ to be a basis of $\mathcal{B}_Q$ forcing ψ_Q to be an isomorphism: due to the conditions (1) and (2), B_W is a submonoid of $\mathcal{B}_Q$ which contains the generators X_q of $\mathcal{B}_Q$ and the unit 1. Hence the k-linear span of B_W is the monoid algebra $\mathcal{B}_Q$ itself. Furthermore condition (3) ensures that the elements of B_W are indexed by W. Thus B_W is a k-linear basis of $\mathcal{B}_Q$ with $|W|$ elements. Moreover the canonical epimorphism ψ_Q is an isomorphism because of condition (3). The existence of an admissible normal form is not obvious. But finding one summarizes our strategy of proving that the relations are defining in several special cases.

3.1 Tools for (2) and reductions

To begin with one needs a better understanding of the multiplication of two arbitrary monomials in $\mathcal{B}_Q$, that is to say of the defining relations. Since Q is acyclic the third relation (under the two first ones) is equivalent to the following two:

- $X_tX_sX_t = X_sX_t$ for all $s, t \in Q_0$, such that there is an arrow $\alpha\colon s \to t$
- $X_sX_t = X_tX_s$ for all vertices $s, t \in Q_0$ which are not connected by an arrow

Therefore the underlying monoid of $\mathcal{B}_Q$ is isomorphic to the Hecke-Kiselman semigroup associated with Q introduced in [GM11]. Let us fix some more notation. We will write $\{v\}$ for the set of the letters occuring in the word $v \in Q_0^*$. For example we have $\{v\} = \{1, 5, 7, 15\}$ if $Q_0 = \{1, 2, \dots, 15\}$ and $v = 1\,.\,5\,.\,15\,.\,7\,.\,1\,.\,5$. For a subset M of Q_0 we will denote by Q_M the full subquiver of Q whose underlying set of vertices is exactly M and we will abbreviate $Q_v := Q_{\{v\}}$. A vertex $t \in Q_0$ is called a sink (source), if no arrow has tail (head) t. The condition on $s, t \in Q_0$ in the third relation defining $\mathcal{B}_Q$ could be replaced by requiring t to be a sink in $Q_{s\,.\,t}$. The defining relations of $\mathcal{B}_Q$ generalize to the following identities in $\mathcal{B}_Q$ (and thus in $k\pi_Q$):

Lemma 3.2. *For all $t \in Q_0$ and all words v, w over Q_0 we have:*

$$X_t X_w X_t = X_w X_t \qquad \text{if } t \text{ is a sink in the subquiver } Q_{t\,.\,w}$$

$$X_t X_w X_t = X_t X_w \qquad \text{if } t \text{ is a source in the subquiver } Q_{t\,.\,w}$$

$$X_v X_w = X_w X_v \qquad \text{if there is no arrow between the subquivers } Q_v \text{ and } Q_w$$

Proof. The second identity follows by duality from the first one, meanwhile the third one results directly from the others. We prove the first identity by an induction on the length of w. If $w = s$ the identity is just one of the defining relations of $\mathcal{B}_Q$. So let $w = u\,\mathbf{.}\,s$ for some $s \in Q_0$ and some word u. The induction hypothesis applies now to s and to u:

$$X_t X_w X_t = X_t X_u X_s X_t = X_t X_u X_t X_s X_t = X_u X_t X_s X_t = X_u X_s X_t = X_w X_t$$

□

Essentially all calculations are abbreviated using these generalized relations. For example the finiteness of π_Q can be deduced from them.

Corollary 3.3. *The k-algebra $\mathcal{B}_Q$ is finite dimensional. Hence the monoid π_Q is finite.*

Proof. If Q just consists of one vertex, $\mathcal{B}_Q$ is two-dimensional. So now we assume Q to have at least two vertices, pick a sink $s \in Q_0$ and consider the quiver $K := Q_{Q_0 \setminus \{s\}}$. Inductively $\mathcal{B}_K \subseteq \mathcal{B}_Q$ is finite dimensional, hence has a finite basis B of monomials over $\{X_t \mid t \in Q_0 \setminus \{s\}\}$. Due to Lemma 3.2 every monomial $X_w \in \mathcal{B}_Q$ lies either in $\mathcal{B}_K X_s \mathcal{B}_K$ (if $s \in \{w\}$) or in $\mathcal{B}_K$ (if $s \notin \{w\}$). Thus $B \cup B X_s B$ is a finite k-linear generating system of $\mathcal{B}_Q$. □

The next observations enable us to restrict to isomorphism classes of finite, acyclic and connected quivers without multiple arrows. We will call a subquiver $K = (K_0, K_1)$ of Q “the quiver reduced by multiple arrows of Q“ if $K_0 = Q_0$ holds and if there is exactly one arrow between two vertices s and t in K_1 whenever there exists (at least) one arrow between s and t in Q_1.

Proposition 3.4. *Let K, K' and Q be finite, acyclic quivers.*
- *(a) If K is a full subquiver of Q, then $\mathcal{B}_K$ is a subalgebra of $\mathcal{B}_Q$.*
- *(b) If Q and K are (anti-) isomorphic, then $\mathcal{B}_Q$ and $\mathcal{B}_K$ are (anti-) isomorphic.*
- *(c) If K is the quiver reduced by multiple arrows of Q, then $\mathcal{B}_K$ and $\mathcal{B}_Q$ are isomorphic.*
- *(d) If K and K' are the connected components of Q, then $\mathcal{B}_Q$ and $\mathcal{B}_K \otimes_k \mathcal{B}_{K'}$ are isomorphic.*

The analogous statements for $k\pi_K$ and $k\pi_Q$ are a priori not clear. However, if there is an admissible normal form associated with K most of them hold (see Proposition 3.6).

3.2 Tools for (3) and a criteria for reductions

For any vertex $t \in Q_0$ we abbreviate $\mathsf{P}_t = \mathsf{P}_t^{(Q)} := \mathsf{P}_{S_t}$ and similar for t_{S_t}. The projection functor P_t on Q-${}_k$rep associated with S_t is easily computed for representations V of Q, since the functor t_{S_t} maps V to that submodule U of the socle $\mathrm{Soc}(V) \subseteq V$, which is given by $U_t = (\mathrm{Soc} V)_t$ and $U_s = 0$ for all $s \in Q_0 \backslash \{t\}$. Therefore $\mathsf{P}_t V$ is described by $(\mathsf{P}_t V)_t = V_t / U_t$ and $(\mathsf{P}_t V)_s = V_s$ for all

$s \in Q_0 \backslash \{t\}$ and the respectively induced k-linear maps. Consequences, in particular for simples and injective indecomposable representations, are summarized in the next remarks. Let I_x be the injective envelope of S_x.

Remark 1. *A Q-representation V is fixed under the action of those P_t with $t \notin \operatorname{supp} V := \{q \in Q_0 \mid V(q) \neq 0\}$. Thus $\mathsf{P}_w V = V$ holds for all words w over $Q_0 \backslash \operatorname{supp}(V)$. In particular we have for all $w \in Q_0^*$:*

$$\mathsf{P}_w(S_t) = \begin{cases} S_t & \text{if } t \notin \{w\} \\ 0 & \text{if } t \in \{w\} \end{cases}$$

Remark 2. *The action of P_t on the injective indecomposable representation I_x is:*

$$\mathsf{P}_t(I_x) = \begin{cases} I_x & \text{if } x \neq t \\ I_t/S_t = \bigoplus_{s \to t} I_s & \text{if } x = t \end{cases}$$

Hence I_x is fixed under P_w for all words w over Q_0 not containing x.

Let K be a subquiver of Q and $F\colon K\text{-}_k\mathrm{rep} \to Q\text{-}_k\mathrm{rep}$ the canonical embedding functor. Recall that for every K-representation U the Q-representation FU is defined by setting for all $t \in Q_0$ and $\alpha \in Q_1$:

$$(FU)_t := \begin{cases} U_t & \text{if } t \in K_0 \\ 0 & \text{otherwise} \end{cases} \quad \text{and} \quad (FU)_\alpha := \begin{cases} U_\alpha & \text{if } \alpha \in K_1 \\ 0 & \text{otherwise} \end{cases}$$

Lemma 3.5. *Let K be a subquiver of Q, $x \in K_0$ and $v, w \in K_0^*$.*
(a) The functors $F\mathsf{P}_x^{(K)}$ und $\mathsf{P}_x^{(Q)}F$ are naturally isomorphic.
(b) If there is a K-representation V with $\mathsf{P}_v^{(K)}V \ncong \mathsf{P}_w^{(K)}V$, then $\mathsf{P}_v^{(Q)}$ and $\mathsf{P}_w^{(Q)}$ are not naturally isomorphic.

Proof. We apply F to the short exact sequence of a K-representation U induced by $\mathsf{P}_x^{(K)}$ and get the short exact sequence:

$$0 \longrightarrow F\mathsf{t}_x^{(K)}U \longrightarrow FU \longrightarrow F\mathsf{P}_x^{(K)}U \longrightarrow 0$$

This is already the short exact sequence of FU induced by $\mathsf{P}_x^{(Q)}$ because of $F\mathsf{t}_x^{(K)}U = \mathsf{t}_x^{(Q)}FU$ and the uniqueness (up to isomorphism) of the cokernel. Therefore (a) holds. In particular, we conclude (b) from: $\mathsf{P}_v^{(Q)}FV \cong F\mathsf{P}_v^{(K)}V \ncong F\mathsf{P}_w^{(K)}V \cong \mathsf{P}_w^{(Q)}FV$. □

Proposition 3.6. *Let K, K' and Q be finite, acyclic quivers. Assume that there are admissible normal forms W and W' associated with K and K' respectively.*
(a) If K is a full subquiver of Q, then $k\pi_K$ is a subalgebra of $k\pi_Q$.
(b) If Q and K are isomorphic, then $k\pi_K$ and $k\pi_Q$ are isomorphic.
(c) If K is the quiver reduced by multiple arrows of Q, then $k\pi_K$ and $k\pi_Q$ are isomorphic.
(d) If K and K' are the connected components of Q, then $k\pi_Q$ and $k\pi_K \otimes_k k\pi_{K'}$ are isomorphic.

Proof. By the assumption on K we have $k\pi_K \cong \mathcal{B}_K$. In case (a) or (c), the assignment $\mathsf{P}_x^{(K)} \mapsto \mathsf{P}_x^{(Q)}$ for all $x \in K_0$ extends to a homomorphism $k\pi_K \longrightarrow k\pi_Q$. Its image is linearly spanned by the monomials $\mathsf{P}_v^{(Q)}$ with $v \in W$. But by the previous lemma, Lemma 3.5, the tuple $(\mathsf{P}_v^{(Q)} | v \in W)$ is also linearly independent. Thus the assertions (a) - (c) hold for dimensional reasons. Meanwhile an admissible normal form associated with Q in case (d) is given by $W \centerdot W'$. Therefore $k\pi_Q \cong \mathcal{B}_Q$ holds and (d) follows with Proposition 3.4(d). ◻

3.3 The linearly oriented Dynkin quiver of type A

We denote by $\mathcal{P}_n$ the poset of the product order $\geq_n$ on the powerset of $\underline{n}\rfloor$ defined in the introduction. For two intervals J and I in $\underline{n}\rfloor$ we define $J \succ I$ by requiring $\min J > \min I$ and $\max J > \max I$, in particular $\{\max J, \max I\} \geq_n \{\min J, \min I\}$. The poset $\mathcal{P}_n$ is in bijection with the set consisting of tuples of intervals $J_r \succ \ldots \succ J_1$ in $\underline{n}\rfloor$. For an interval $J = \{i, i+1, \ldots, j-1, j\}$ of positive integers let J denote the word $i \centerdot i+1 \centerdot \ldots \centerdot j-1 \centerdot j$ as well. The monomial $X_J \in \mathcal{B}_{Q_n}$ is an idempotent. More precisely for all $k \in J$ we have $X_J X_k = X_J$ since k is a source in the subquiver $Q_{k \centerdot k+1 \centerdot \ldots \centerdot j-1 \centerdot j}$ (Lemma 3.2). We will meet a generalisation of these idempotents to arbitrary finite quivers without oriented cycles to determine the radical of $\mathcal{B}_Q$ in the next section.

Proposition 3.7. *An admissible normal form of $\mathcal{B}_{Q_n}$ is*

$$W_n := \{J_r \centerdot \ldots \centerdot J_1 \mid r \in \underline{n}\rfloor \cup \{0\},\ \ J_r \succ \ldots \succ J_1 \ \textit{ intervals in } \underline{n}\rfloor\}$$

So $\mathcal{B}_{Q_n}$ and $k\pi_{Q_n}$ are isomorphic and have the dimension $C_{n+1} = \frac{1}{n+2}\binom{2(n+1)}{n+1}$ by characterization 6.19.aa. in [Sta99] of the $n+1^{\text{th}}$ Catalan number C_{n+1}.

Proof. Condition (1) is easily verified by looking at $r = 0$ and $r = 1$. To check condition (2) we just state a multiplication rule:

Let $n \in \mathbb{N}$. For all intervals J and $L_1 \prec \ldots \prec L_s$ in $\underline{n}\rfloor$ and for $L_0 := \emptyset =: L_{s+1}$ we define indices $y = y(J, L_1, \ldots, L_s)$ and $z = z(J, L_1, \ldots, L_s)$ by:

$$y := \begin{cases} \max\{x \in \underline{s}\rfloor \mid J \succ L_x \textit{ and } J \cap (L_x + 1) = \emptyset\} & \textit{if } L_1 \prec J \textit{ and } J \cap (L_1 + 1) = \emptyset \\ 0 & \textit{else} \end{cases}$$

and

$$z := \begin{cases} y + 1 & \textit{if } J \prec L_{y+1} \\ \min\{x \in \{y+2, \ldots, s\} \mid L_{y+1} \cup J \prec L_x\} & \textit{if } J \nprec L_{y+1} \textit{ and } J \cup L_{y+1} \prec L_s \\ s + 1 & \textit{else, thus if } J \nprec L_{y+1} \textit{ and } J \cup (L_{y+1}) \nprec L_s \end{cases}$$

Then the product of $X_{L_1 \ldots L_s}$ and X_J is given by:

$$X_{L_s \ldots L_1} X_J = X_{L_s \ldots L_z (J \cup \bigcup_{t=y+1}^{z-1} L_t) L_y \ldots L_1} = \begin{cases} X_{L_s \ldots L_{y+1} J L_y \ldots L_1} & \textit{if } J \prec L_{y+1} \\ X_{L_s \ldots L_z (J \cup L_{y+1}) L_y \ldots L_1} & \textit{otherwise} \end{cases}$$

The proof is a lengthy but straightforward induction on s requiring case by case analysis. Now we turn towards condition (3). For this it suffices to consider the injective indecomposable Q_n-representations I_j for $j \in \underline{n}\rfloor \cup \{0\}$. We set $I_0 := 0$. Because of $\mathsf{P}_j I_j = I_{j-1}$ we get inductively on r for all intervals $J_r \succ \ldots \succ J_1$ in $\underline{n}\rfloor$:

$$\mathsf{P}_{J_r \ldots J_1} I_j = \begin{cases} I_{\min J_a - 1} & \text{if } j \in \bigcup_{x=1}^{r} J_x, \text{ where } a \in \underline{r}\rfloor \text{ is min. with } j \in J_a \\ I_j & \text{otherwise} \end{cases}$$

Let $J_r \succ \ldots \succ J_1$ and $L_s \succ \ldots \succ L_1$ be two distinct tuples of intervals in $\underline{n}\rfloor$. Without loss of generality we can pick an index $a \in \underline{r}\rfloor$ minimal with respect to J_a being distinct from $L_1, \ldots, L_s$. Then we have:

$$\mathsf{P}_{J_r \ldots J_1}(I_{\max J_a}) = I_{\min J_a - 1}$$

and

$$\mathsf{P}_{L_s \ldots L_1}(I_{\max J_a}) = \begin{cases} I_{\min L_b - 1} & \text{if } \max J_a \in \bigcup_{x=1}^{s} L_x, \text{ where } b \in \underline{s}\rfloor \text{ is min. with } \max J_a \in L_b \\ I_{\max J_a} & \text{otherwise} \end{cases}$$

We are done if $\max J_a \notin \bigcup_{x=1}^{s} L_x$. So we assume $\max J_a \in \bigcup_{x=1}^{s} L_x$. We are done as well if $\min L_b \neq \min J_a$. Thus let $\min L_b = \min J_a$. Now we consider the action on $I_{\max L_b}$:

$$\mathsf{P}_{L_s \ldots L_1} I_{\max L_b} = I_{\min L_b - 1} = I_{\min J_a - 1}$$

and

$$\mathsf{P}_{J_1 \ldots J_r} I_{\max L_b} = \begin{cases} I_{\min J_c - 1} & \text{if } \max L_b \in \bigcup_{x=1}^{r} J_x, \text{ where } c \in \underline{r}\rfloor \text{ is min. with } \max L_b \in J_c \\ I_{\max L_b} & \text{otherwise} \end{cases}$$

In the first case (for $\max L_b$) the inequality holds because of $\max J_c \geq \max L_b > \max J_a$, so $J_c \succ J_a$, hence $\min J_c > \min J_a$. In the second one it holds due to $\max L_b > \min L_b - 1$. □

The defining relations for π_{Q_n} are the same as for the monoid NDPF_{n+1} of non-decreasing parking functions (see [HT09] or [GM11]) which is generated by the functions $\pi_j := \begin{pmatrix} 1 \ldots j-1 & j & j+1 \ldots n+1 \\ 1 \ldots j-1 & j & j \quad \ldots n+1 \end{pmatrix}$. In fact, if β is the bijection $\{I_0, I_1, \ldots, I_n\} \longrightarrow \underline{n+1}\rfloor, I_j \mapsto j+1$ the proof of condition (3) yields the isomorphism $\pi_{Q_n} \longrightarrow \mathrm{NDPF}_{n+1}, \mathsf{P}_v \mapsto \beta \mathsf{P}_v \beta^{-1}$, which is the extension of the assignment $\mathsf{P}_j \mapsto \pi_j$. Thus the full subcategory of the category of covariant functors on kQ-mod containing the elements $\mathsf{P}_{t_1} \mathsf{P}_{t_2} \ldots \mathsf{P}_{t_r}$ for $r \geq 0$ and vertices $t_1, t_2, \ldots, t_r \in Q_0$ categorifies the monoid NDPF_{n+1}.

3.4 Gluing on a sink

We start with n finite acyclic and pairwise disjoint quivers $Q(1), \ldots, Q(n)$ and pick some vertices $p_1^{(1)}, \ldots, p_{r_1}^{(1)} \in Q(1), \ldots, p_1^{(n)}, \ldots, p_{r_n}^{(n)} \in Q(n)$. Then we consider the quiver Q arising from gluing these quivers together on a new vertex s over new arrows $\alpha_i^{(j)} \colon p_i^{(j)} \to s$. The shape of

Q is sketched below. We will moreover denote by $\widetilde{\mathrm{Q}}(j)$ the full subquiver of Q with the vertices $Q(j)_0 \cup \{s\}$.



Lemma 3.8. *Assume that for every* $j \in \underline{n}\rfloor$ *we have admissible normal forms* $\mathrm{W}(j) \subseteq Q(j)_0^*$ *and* $\widetilde{\mathrm{W}}(j) \subseteq \widetilde{\mathrm{Q}}(j)_0^*$ *associated with* $Q(j)$ *and* $\widetilde{\mathrm{Q}}(j)$ *respectively, such that* $\mathrm{W}(j) \subseteq \widetilde{\mathrm{W}}(j)$ *holds and* s *appears at most once in any word of* $\widetilde{\mathrm{W}}(j)$*. Then an admissible normal form* W *associated with* Q *consists of the* $\prod_{j\in\underline{n}\rfloor} |\mathrm{W}(j)| + \prod_{j\in\underline{n}\rfloor}(|\widetilde{\mathrm{W}}(j)| - |\mathrm{W}(j)|)$ *words:*

$$(a) \qquad w_1 \centerdot w_2 \centerdot \ldots \centerdot w_n \qquad \textit{with } \ w_j \in \mathrm{W}(j) \ \textit{ for all } \ j \in \underline{n}\rfloor$$

and

$$(b) \qquad y_1 \centerdot y_2 \centerdot \ldots \centerdot y_n \centerdot s \centerdot z_1 \centerdot z_2 \centerdot \ldots \centerdot z_n \qquad \textit{with } \ y_j \centerdot s \centerdot z_j \in \widetilde{\mathrm{W}}(j) \setminus \mathrm{W}(j) \ \textit{ for all } \ j \in \underline{n}\rfloor$$

Thus $\mathcal{B}_Q \cong k\pi_Q$ *holds. Moreover the dual assertions holds.*

Proof. Conditions (1) and (2) are verified straightforwardly using the generalized relations. For condition (3) it suffices, by the remarks in subsection 3.2, to consider two words $v \neq w \in W$ both being either of type (a) or of type (b). In each case there are an index $j \in \underline{n}\rfloor$ and subwords $v_j, w_j \in \widetilde{\mathrm{W}}(j)$ of v and w respectively such that $v_j \neq w_j$ holds. Now employ the assumption on $\widetilde{\mathrm{W}}(j)$ and Lemma 3.5. □

A direct application of this lemma gives an admissible normal form associated with the m-subspace quiver T_m, which is the connected quiver with exactly one sink s and m sources enumerated by $1, \ldots, m$: here $Q(j)$ corresponds just to the vertex j and $\widetilde{\mathrm{Q}}(j)$ to $j \to s$. An admissible normal form associated to the latter is $\{\emptyset, j, s, js, sj\}$ which contains the normal form associated with j, i.e. $\{\emptyset, j\}$. To fix an order on the vertices we denote for every subset $J = j_1 < \ldots < j_k$ of $\underline{m}\rfloor$ the word $j_1 \centerdot \ldots \centerdot j_k$ with $w(J)$. Now an admissible normal form associated with T_m has $2^m + 3^m$ elements:

$$\{w(J) \mid J \subseteq \underline{m}\rfloor\} \ \cup \ \{w(I) \centerdot s \centerdot w(J) \mid I, J \subseteq \underline{m}\rfloor, I \cap J = \emptyset\}$$

This can be extended to the star quiver, since we have admissible normal forms associated with its branches, i.e linearly oriented Dynkin quivers of type A.

3.4.1 Tree quivers with a specific orientation

Furthermore, Lemma 3.8 is the induction step for tree quivers with a specific orientation: we will call Q an admissible tree quiver, if each crossing of Q, i.e. a vertex whose entry degree or exit degree is at least 2, is either a sink or a source. (So the linearly oriented Dynkin quivers D_4 are not subquivers of Q.) One can endow every tree with an orientation to obtain an admissible tree quiver. In particular every tree with a bipartite orientation is an admissible tree quiver. Note also that every Dynkin quiver of type A is an admissible tree quiver.

Theorem 3.9. *There is an admissible normal form associated with any admissible tree quiver Q. In particular the relations are defining for π_Q.*

Proof. The case Q being Q_n for some n is already done. So assume Q not to be a linear oriented Dynkin quiver of type A. In particular we can pick a crossing s of Q. To apply Lemma 3.8 we just have to check, whether the assumptions hold for the subquivers which are linked to s by one arrow. These subquivers (and their extensions with s) are again admissible tree quivers. So it suffices to show:
Let K be an admissible tree quiver, $y \in K_0$ and $\widetilde{\mathrm{K}}$ an extension of K by one (new) vertex s and one (new) arrow $y \leftarrow s$ or $y \rightarrow s$, so that $\widetilde{\mathrm{K}}$ is again an admissible tree quiver. Then there are admissible normal forms $\mathrm{W} \subseteq K_0^*$ and $\widetilde{\mathrm{W}} \subseteq \widetilde{\mathrm{K}}_0^*$ associated with K and $\widetilde{\mathrm{K}}$ respectively, such that $\mathrm{W} \subseteq \widetilde{\mathrm{W}}$ and s appears at most once in any word of $\widetilde{\mathrm{W}}$.
This is proven by induction on the number of vertices of K using Lemma 3.8. □

The induction step, i.e. Lemma 3.8, provides a procedure for gaining an admissible normal form associated with Q. We illustrate this by the special case of bipartite Dynkin quivers K_n of type A_n. Depending on whether n is even or odd K_n has up to anti-isomorphism one of the following shapes:



Admissable normal forms $\mathrm{W}_1 \subseteq \mathrm{W}_2 \subseteq \mathrm{W}_3$ associated with $K_1 \subseteq K_2 \subseteq K_3$ respectively are:

$$\{\emptyset, 1\} \subseteq \{\emptyset, 1, 2, 12, 21\} \subseteq \{\emptyset, 1, 3, 13, 2, 12, 21, 32, 23, 123, 321, 132, 213\}$$

For $n \geq 4$ the vertex $n-1$ is a crossing and either a sink or a source. Inductively we have admissible normal forms $\mathrm{W}_{n-2} \subseteq \mathrm{W}_{n-1}$ associated with $K_{n-2} \subseteq K_{n-1}$ respectively such that $n-1$ appears at most once in any word of W_{n-1}. On the other hand we have admissible normal forms $\{\emptyset, n\} \subseteq \{\emptyset, n, n-1, n \mathbf{.} n-1, n-1 \mathbf{.} n\}$ associated with the quivers n and $n-1 \rightarrow n$ (or $n-1 \leftarrow n$) respectively. Now the Lemma 3.8 with $s = n-1$ yields the admissible normal form

$$\mathrm{W}_n := \mathrm{W}_{n-2} \cup \mathrm{W}_{n-2} \mathbf{.} n \bigcup \mathrm{W}_{n-1} \setminus \mathrm{W}_{n-2} \bigcup n \mathbf{.} (\mathrm{W}_{n-1} \setminus \mathrm{W}_{n-2}) \bigcup (\mathrm{W}_{n-1} \setminus \mathrm{W}_{n-2}) \mathbf{.} n$$

which contains W_{n-1} and fulfils the condition on the appearance of n. Therefore the dimension $|\mathrm{W}_n|$ of $k\pi_{K_n} \cong \mathcal{B}_{K_n}$ can be calculated over the recurrence relation:

$$|\mathrm{W}_n| = 2|W_{n-2}| + 3(|W_{n-1}| - |W_{n-2}|) = 3|W_{n-1}| - |W_{n-2}|$$

Hence $(|\mathrm{W}_j|)_{j \in \mathbb{N}}$ corresponds to the partial sequence $(F_{2j+1})_{j \in \mathbb{N}}$ of the Fibonacci-sequence $(F_n)_{n \in \mathbb{N}}$ with $F_1 = F_2 = 1$.

4 The Gabriel Quiver of $\mathcal{B}_Q$

As before let k be a field and Q a finite, acyclic quiver. Theorem 1.2 is a direct consequence of the next two subsections.
One remark before we start. In [DHST11] the authors determined among other things the radical and Gabriel quiver of the monoid algebra of a finite $\mathcal{J}$-trivial monoid. This could be applied here, because $\mathcal{B}_Q$ is – as the monoid algebra of the Hecke-Kiselman monoid HK_Q – such a monoid algebra. We repeat the argument mentioned in [GM11]: Since being $\mathcal{J}$-trivial is closed under quotients (see [Pin]), it suffices to show, that HK_Q is a quotient of a $\mathcal{J}$-trivial monoid. Now the Hecke-Kiselman monoid HK_{K_n} associated to the quiver K_n with n vertices $\{1,\dots,n\}$ and arrows $i \to j$ for each pair $i < j$ is the Kiselman semigroup and $\mathcal{J}$-trivial by [KM09]. Moreover Q can be embedded in the quiver K_n for $n := |Q_0|$ by choosing an enumeration $\{1,\dots,n\}$ of the vertices Q_0 such that $i \to j$ implies $i < j$. Thus there is the canonical projection introduced in [GM11] from HK_{K_n} onto HK_Q.
Here, we compute the radical and the Gabriel-quiver of $\mathcal{B}_Q$ directly just using the defining relations.

4.1 The simples and the radical of $\mathcal{B}_Q$

The structure of the simple modules are closely related to those of the 0-Hecke algebra (see [Nor79] and [HT09]). For every subset M of Q_0 we define $\mathrm{E}_M = (k, \delta_M)$ to be the (one-dimensional) $\mathcal{B}_Q$-module given by the homomorphism $\delta_M\colon \mathcal{B}_Q \longrightarrow k \cong \mathrm{End}_k(k)$ of algebras with $X_q \mapsto 1$ if $q \in M$ and $X_q \mapsto 0$ otherwise for all $q \in Q_0$. In the sequel we compute the radical to show that this family $(\mathrm{E}_M)_{M\subseteq Q_0}$ of $2^{|Q_0|}$ simple modules represents all simple $\mathcal{B}_Q$-modules. For this we construct for each subset M of Q_0 a specific monomial $X_M \in \mathcal{B}_Q$ to describe a k-linear generating set of the radical. To this end we consider the inductively defined sets of sinks $\mathrm{S}_j M$ at level $j \in \mathbb{N}$ associated to M:

$$\begin{aligned}
\mathrm{S}_0 M &:= \{q \in M \mid q \text{ is a sink in } Q_M\}\\
&\vdots\\
\mathrm{S}_{j+1} M &:= \{q \in M \mid q \text{ is a sink in } Q_{M\setminus(\mathrm{S}_0 M \cup \mathrm{S}_1 M \cup \ldots \cup \mathrm{S}_j M)}\}
\end{aligned}$$

Since M is finite, there is a uniquely determined index $s(M) := m$ such that $\mathrm{S}_m M \neq \emptyset = \mathrm{S}_{m+1} M$ holds. Moreover $Q_{\mathrm{S}_j M}$ contains no arrows. Hence all X_p and X_q with $p, q \in \mathrm{S}_j M$ commute and we can thus define:

$$X_M := \prod_{q\in \mathrm{S}_m M} X_q \;\ldots\; \prod_{q\in \mathrm{S}_0 M} X_q$$

Note that by the generalized relations X_M is idempotent, since we have $X_q X_M = X_M = X_M X_q$ for every $q \in M$.

Proposition 4.1. *The radical* $\mathrm{rad}(\mathcal{B}_Q)$ *of* $\mathcal{B}_Q$ *is the k-linear span of* $\mathcal{M} := \{X_{\{w\}} - X_w \mid w \in Q_0^*\}$. *So* $\mathcal{B}_Q$ *is a basic algebra and* $(\mathrm{E}_M)_{M\subseteq Q_0}$ *is a representative system of its simple modules.*

Proof. Let $\mathcal{V}$ be the k-linear span of $\mathcal{M}$ and $\mathcal{I}$ be the intersection of the annihilators $\mathrm{Ann}_{\mathcal{B}_Q}(\mathrm{E}_M) := \{a \in \mathcal{B}_Q \mid \delta_M(a) = 0\}$ with $M \subseteq Q_0$. Firstly we observe that $\mathcal{V}$ coincides with the ideal $\mathcal{I}$: by the definition of δ_M we have for an arbitrary element $b = \sum_{v\in Q_0^*} b_v X_v$ in $\mathcal{B}_Q$:

$$\delta_M(b) = \sum\nolimits_{v\in Q_0^*,\ \{v\}\subseteq M} b_v$$

Thus b lies in $\mathcal{I}$ if and only if $\sum_{v\in Q_0^*,\, M=\{v\}} b_v = 0$ for all $M \subseteq Q_0$, which is in turn equivalent to

$$b = \sum\nolimits_{M\subseteq Q_0} \sum\nolimits_{v\in Q_0^*, M=\{v\}} b_v(X_v - X_M) \in \mathcal{V}$$

Secondly we show, that $\mathcal{M}$ consists of nilpotent elements. (Hence $\mathcal{V}$ is a nilpotent ideal by a theorem of Wedderburn (see [Pie82], 4.6)). For this we prove the equality $X_w^s = X_{\{w\}}$ for each word w over Q_0 and $s := s(\{w\})$ by an induction on s: if all the letters occuring in w correspond to sinks, X_w already coincides in $\mathcal{B}_Q$ with $X_{\{w\}}$. So now assume $\mathrm{S}_1\{w\} \neq \emptyset$. Furthermore let v be the subword of w, which arises from w by canceling all sinks $x \in \mathrm{S}_0\{w\}$ in w. Since $s(\{v\}) = s - 1$, it follows inductively $X_w^s = X_v^{s-1} X_w = X_{\{v\}} X_w = X_{\{w\}}$ by the generalized relations and the properties of $X_{\{v\}}$ respective $X_{\{w\}}$. Therefore the element $X_{\{w\}} - X_w \in \mathcal{B}_Q$ is nilpotent:

$$(X_{\{w\}} - X_w)^s = (-X_{\{w\}} + X_w^2)(X_{\{w\}} - X_w)^{s-2} = \ldots = (-1)^{s-1}(X_{\{w\}} - X_w^s) = 0$$

Now $\mathcal{V} \subseteq \mathrm{rad}(\mathcal{B}_Q) \subseteq \mathcal{I}$ follows from the different characterizations of the radical of a finite dimensional algebra. □

4.2 The Gabriel quiver of $\mathcal{B}_\mathbf{Q}$

We calculate the k-dimensions of the extension groups $\mathrm{Ext}^1_{\mathcal{B}_Q}(\mathrm{E}_M, \mathrm{E}_N)$, i.e.
the number of arrows from $[\mathrm{E}_M]$ to $[\mathrm{E}_N]$ in the Gabriel quiver $\Gamma(\mathcal{B}_Q)$ of $\mathcal{B}_Q$. Then the algebra $\mathcal{B}_Q$ is a quotient of the path algebra $k\Gamma(\mathcal{B})$ by an ideal I with $\mathrm{rad}(\mathcal{B}_Q)^2 \subseteq I \subseteq \mathrm{rad}(\mathcal{B}_Q)^r$ for some $r \in \mathbb{N}$. We will see that this ideal is zero for the m-subspace quiver as well as for some simply shaped quivers. We devote section 5 to the proof that I is generated by the commutativity relations of $\Gamma(\mathcal{B}_Q)$, if $Q = Q_n$ is the linearly oriented Dynkin quiver of type A.
Since the simple modules are one dimensional, it suffices to determine the two dimensional $\mathcal{B}_Q$-modules. The calculations are similar to those for the 0-Hecke-algebras (type A) – as done for example in [Fay05] with the difference that we have to respect the non-symmetry of the defining relations and the generalisation to finite, acyclic quivers.
Let M and N be subsets of Q_0. We consider the characteristic tuples $(m_q := \delta_M(X_q))_{q\in Q_o}$ and $(n_q := \delta_N(X_q))_{q\in Q_0}$ over $0, 1 \in k$. Then we call a tuple $a = (a_q)_{q\in Q_0}$ over k (or a function $a\colon Q_0 \longrightarrow k, q \mapsto a_q$) admissible or (M, N)-admissible, if the assignment

$$X_q \mapsto \begin{pmatrix} n_q & a_q \\ 0 & m_q \end{pmatrix} =: A_q \in k^{2\times 2}$$

extends uniquely to an homomorphism from $\mathcal{B}_Q$ to $k^{2\times 2}$. We receive a two-dimensional $\mathcal{B}_Q$-module $W(a, M, N) = W(a)$ and up to equivalence the short exact sequences η in $\mathrm{Ext}^1_A(\mathrm{E}_M, \mathrm{E}_N)$ are the sequences η_a:

$$\eta_a : \qquad 0 \longrightarrow \mathrm{E}_N \xrightarrow{\begin{pmatrix}1\\0\end{pmatrix}} W(a) \xrightarrow{\begin{pmatrix}0 & 1\end{pmatrix}} \mathrm{E}_M \longrightarrow 0$$

The dimension of $\operatorname{Ext}^1_{\mathcal{B}_Q}(\mathrm{E}_M, \mathrm{E}_N)$ could be described in terms of the following property for the set-theoretic differences $M\setminus N$ and $N\setminus M$:

Definition 4.2. *Let P and R be two disjoint subsets of Q_0. We say that "P is strongly connected towards R" and write $P \rightrightarrows R$, if neither P nor R are empty and for every two vertices $p \in P$ and $r \in R$ there exists (at least) one arrow from p to r (in Q_1).* [1]

If P is not strongly connected towards R we write $P \not\rightrightarrows R$. Since Q is acyclic, $P \not\rightrightarrows R$ holds if and only if $P = \emptyset$, $R = \emptyset$ or there are vertices $p \in P$ and $r \in R$, such that p is a sink in $Q_{p,r}$.

Lemma 4.3. *(a) We have $\operatorname{Ext}^1_{\mathcal{B}_Q}(\mathrm{E}_M, \mathrm{E}_N) = 0$ if and only if $M\setminus N \not\rightrightarrows N\setminus M$ holds.*
(b) We have $\dim_k \operatorname{Ext}^1_{\mathcal{B}_Q}(\mathrm{E}_M, \mathrm{E}_N) = 1$ if and only if $M\setminus N \rightrightarrows N\setminus M$ holds.

Proof. Firstly we determine the (M,N)-admissible functions. Let $a\colon Q_0 \longrightarrow k, q \mapsto a_q$ be a function. The shape of the assigned A_q depends on whether $q \in M \cap N$, $q \in Q_0 \setminus (M \cup N)$, $q \in M\setminus N$ or $q \in N\setminus M$ and is as follows

$$A_q = \begin{pmatrix}1 & a_q\\0 & 1\end{pmatrix}, \quad A_q = \begin{pmatrix}0 & a_q\\0 & 0\end{pmatrix}, \quad A_q = \begin{pmatrix}0 & a_q\\0 & 1\end{pmatrix}, \quad A_q = \begin{pmatrix}1 & a_q\\0 & 0\end{pmatrix}$$

respectively. If a is an admissable function, then all these matrices are idempotent, i.e. $a_q = 0$ for all $q \in M \cap N$ and for all $q \in Q_0 \setminus (M \cup N)$. Therefore every (M,N)-admissible function lies in

$$\mathcal{F} := \{a\colon Q_0 \to k \mid a|_{M\cap N} = 0 \ \text{ and } \ a|_{Q_0\setminus(M\cup N)} = 0\}$$

and thus it suffices to consider the remaining relations, as for example $A_pA_qA_p = A_qA_pA_q$, just for all $p, q \in M\setminus N \cup N\setminus M$. For example for any two elements p, q in $M\setminus N$ we conclude $a_p = a_q$ from the conditions, that A_pA_q equals one of the products $A_qA_pA_q$ or A_qA_p. Similar considerations show that if $M\setminus N \rightrightarrows N\setminus M$ holds then the set of all admissible functions is

$$\{a \in \mathcal{F} \mid \exists\, c_M, c_N \in k : \quad a|_{M\setminus N} = c_M \operatorname{id}_{M\setminus N} \ \text{ and } \ a|_{N\setminus M} = c_N \operatorname{id}_{N\setminus M}\}$$

Whereas in the case $M\setminus N \not\rightrightarrows N\setminus M$ the existence of $M\setminus N \ni q \leftarrow p \in N\setminus M$ yields the equality $A_pA_q = A_pA_qA_p$, i.e. $a_q + a_p = 0$, hence the set of admissible functions is

$$\{a \in \mathcal{F} \mid \exists c \in k : \quad a|_{M\setminus N} = c \operatorname{id}_{M\setminus N} \ \wedge \ a|_{N\setminus M} = -c \operatorname{id}_{N\setminus M}\} = \langle (m_q - n_q)_{q\in Q_0} \rangle_k$$

So $M\setminus N \not\rightrightarrows N\setminus M$ holds if and only if every (M,N)-admissible function lies in $\langle (m_q - n_q)_{q\in Q_0} \rangle_k$. Hence it suffices to show that $\operatorname{Ext}^1_{\mathcal{B}_Q}(\mathrm{E}_M, \mathrm{E}_N) = 0$ holds iff every (M,N)-admissible function lies in $\langle (m_q - n_q)_{q\in Q_0} \rangle_k$. For this let η_0 denote the trivial short exact sequence and $W(0)$ its middle term; in particular $W(0) \cong \mathrm{E}_M \oplus \mathrm{E}_N$. We show for every admissible function a the equivalence:

$$(*) \quad \eta_a \sim \eta_0 \iff a \in \langle (m_q - n_q)_{q\in Q_0} \rangle_k$$

---

[1] In this case the subquiver of Q with the vertices $P \cup R$ and just those arrows from Q_1, which connect vertices from P to vertices from R, is a completely bipartite quiver.

If $c \in k$ and $(a_q = cm_q - cn_q)_{q \in Q_0}$ an admissible function, then a $\mathcal{B}_Q$-homomorphism from $W(a)$ to $W(0)$, which additionally provides $\eta_a \sim \eta_0$, is given by the left multiplication with $\begin{pmatrix} 1 & c \\ 0 & 1 \end{pmatrix}$. On the other hand every homomorphism $\Phi \colon W(a) \longrightarrow W(0)$ providing $\eta_a \sim \eta_0$ is given by a matrix $\begin{pmatrix} 1 & c \\ 0 & 1 \end{pmatrix}$ for a constant $c \in k$ by the commutativity of the corresponding diagram. Since Φ is in particular a $\mathcal{B}_Q$-homomorphism, it follows $a_p + cm_p = cn_p$ for all $p \in Q_0$. Hence (a) is proved. Statement (b) follows from (a) if one shows that any two non-trivial short exact sequences are linearly dependent. So let a and b be (M, N)-admissible functions such that $[\eta_0] \neq [\eta_a] \neq [\eta_b] \neq [\eta_0]$ holds. By $(*)$ and the previous thoughts there exist $c \neq c'$ resp. $d \neq d'$ in k with

$$a|_{M\backslash N} = c\,\mathrm{id}_{M\backslash N} \quad \text{and} \quad a|_{N\backslash M} = c'\,\mathrm{id}_{N\backslash M} \quad \text{resp.} \quad b|_{M\backslash N} = d\,\mathrm{id}_{M\backslash N} \quad \text{and} \quad b|_{N\backslash M} = d'\,\mathrm{id}_{N\backslash M}$$

Then $e := d + d'/c + c' \neq 0$ and the left multiplication with $\begin{pmatrix} e & c'e - d' \\ 0 & 1 \end{pmatrix}$ is a $\mathcal{B}_Q$-isomorphism from $W(a)$ to $W(b)$ providing

$$\eta_a \sim e^{-1}\eta_b \colon 0 \longrightarrow \mathrm{E}_N \xrightarrow{\begin{pmatrix} e & 0 \end{pmatrix}} W(b) \xrightarrow{\begin{pmatrix} 0 \\ 1 \end{pmatrix}} \mathrm{E}_M \longrightarrow 0$$

□

4.3 Properties of the Gabriel quiver

Let Q be a finite acyclic quiver. For this section we abbreviate notation: instead of $[\mathrm{E}_M]$ we just write M. Accordingly, the set of vertices of the Gabriel quiver is from now on just the powerset $\wp(Q_0)$ of Q_0 . In this notation there is (exactly) one arrow from $M \in \wp(Q_0)$ to $N \in \wp(Q_0)$ in $\Gamma(\mathcal{B}_Q)$, if $M\backslash N$ is strongly connected towards $N\backslash M$ (w.r.t. Q). In particular, the Gabriel quiver has no loops. Obviously, Q can be embedded in $\Gamma(\mathcal{B}_Q)$. Besides Q^{op} is isomorphic to the full subquiver of $\Gamma(\mathcal{B}_Q)$ with the vertices $\{Q_0\backslash\{p\} \mid p \in Q_0\}$. In general, we have for all M and N:

$$M \to N \in \Gamma(\mathcal{B}_Q) \Longleftrightarrow Q_0\backslash N \to Q_0\backslash M \in \Gamma(\mathcal{B}_Q)$$

Therefore the map $\wp(Q_0) \longrightarrow \wp(Q_0)$, $M \mapsto Q_0\backslash M$ induces an involution ι on $\Gamma(\mathcal{B}_Q)$. In particular we have in the case that $\mathcal{B}_Q$ is already isomorphic to $k\Gamma(\mathcal{B}_Q)$: $\mathcal{B}_Q \cong \mathcal{B}_Q^{op} \cong \mathcal{B}_{Q^{op}}$. Now we look at special cases and some examples and end this chapter with general observations.

- The Gabriel quiver of $\mathcal{B}_{Q_n}$ is described by the following equivalence:

  There is one arrow $M \to N$ in $\Gamma(\mathcal{B}_{Q_n})$ if and only if there exists exactly one index $i \in \underline{n\rfloor}$ with $M\backslash N = \{i\}$ and $N\backslash M = \{i+1\}$.

  Therefore only equally large sets are connected in $\Gamma(\mathcal{B}_{Q_n})$. Thus the Gabriel quiver of $\mathcal{B}_{Q_n}$ has at least $n+1$ connected components. Actually $\Gamma(\mathcal{B}_Q)$ has exactly $n+1$ as we will see.

- In the m-subspace quiver T_m each subset of sources is strongly connected towards the only sink set $\{s\}$. Thus we have:

  There is one arrow $M \to N$ in $\Gamma(\mathcal{B}_Q)$ if and only if there exist two disjoint subsets $M' \neq \emptyset$ and N' of $\underline{m\rfloor}$ with $M = N' \cup M'$ and $N = N' \cup \{s\}$.

  As in the first special case $\emptyset$ and $(T_m)_0$ are isolated vertices of the Gabriel quiver. Moreover each vertex $\emptyset \neq M \neq (T_m)_0 \in \wp(Q_0)$ is either a source (if $M \subseteq \underline{m\rfloor}$) or a sink (if $s \in M$). Hence there are no paths of length ≥ 2. Consequently the algebra $k\Gamma(\mathcal{B}_Q)$ has radical square 0 and is already isomorphic to $\mathcal{B}_Q$. In particular $\mathcal{B}_Q \cong \mathcal{B}_Q^{op}$. By the way the only arrow in $\Gamma(\mathcal{B}_Q)$ which is fixed under the above presented involution ι is $\underline{n\rfloor} \to \{s\}$.

We assume Q to be connected. We call $q \in Q_0$ a successor of $p \in Q_0$, if there is a path in Q from p to q. By Theorem 1.2 it follows straightforwardly that the sinks (sources) of the Gabriel quiver of $\mathcal{B}_Q$ are precisely the subsets of Q_0 which are closed under successors (predecessor) in Q. Therefore we now know the projective (injectives) amongst the simple $\mathcal{B}_Q$-modules. We end this section with the proof of Proposition 1.3. Successively applying the next lemma shows that the full subquiver $\Gamma(\mathcal{B}_Q)_j$ of $\Gamma(\mathcal{B}_Q)$ whose vertices are those subsets of Q_0 with exactly j elements is connected for every $j \leq |Q_0|$:

Lemma 4.4. *For every proper non-empty subset A of Q_0, each $a \in A$ and each $b \in Q_0 \backslash A$ there exists a walk between A and $A\backslash\{a\} \cup \{b\}$ in $\Gamma(\mathcal{B}_Q)$.*

Proof. In the sequel we write $x -\!\!- y$, if the vertices x and y are connected by an arrow. Since Q is connected, there is a walk between any two vertices a and b in Q_0:

$$a = x_0 -\!\!- x_1 -\!\!- \ldots -\!\!- x_r -\!\!- x_{r+1} = b$$

Without loss of generality we can assume, that $x_0, \ldots, x_{r+1}$ are pairwise disjoint. Based on such a walk we construct inductively on the number of the changes from A to $Q_0 \backslash A$, i.e. on the number n of indices $j \in \underline{r+1\rfloor} \cup \{0\}$ with $x_j \in A$ and $x_{j+1} \in Q_0 \backslash A$, a walk in $\Gamma(\mathcal{B}_Q)$ between A and $A\backslash\{a\} \cup \{b\}$.

$\underline{n = 1}$: Let $a \in A \subseteq Q_0$ and $b \in Q_0 \backslash A$, such that there is a walk in Q of the following kind:

$$\underbrace{a -\!\!- x_1 -\!\!- \ldots -\!\!- x_{j-1}}_{\in A} -\!\!- \underbrace{x_j -\!\!- \ldots -\!\!- x_r -\!\!- b}_{\in Q_0 \backslash A}$$

Then we obtain the following walk in the Gabriel quiver by Theorem 1.2:

$$\begin{aligned} &A -\!\!- A\backslash\{x_{j-1}\} \cup \{x_j\} -\!\!- A\backslash\{x_{j-1}\} \cup \{x_{j+1}\} -\!\!- \ldots -\!\!- A\backslash\{x_{j-1}\} \cup \{x_r\} -\!\!- A\backslash\{x_{j-1}\} \cup \{b\} \\ &\quad -\!\!- A\backslash\{x_{j-2}\} \cup \{b\} -\!\!- \ldots -\!\!- A\backslash\{x_1\} \cup \{b\} -\!\!- A\backslash\{a\} \cup \{b\} \end{aligned}$$

$\underline{n \to n+1}$: Now let $a \in A \subseteq Q_0$ und $b \in Q_0 \backslash A$ such that, there is a walk in Q over pairwise disjoint vertices between a and b with more than one change between A and $Q_0 \backslash A$. Such a walk is of the kind:

$$\underbrace{a = x_0 -\!\!- \ldots -\!\!- x_{j-1}}_{\in A} -\!\!- \underbrace{x_j -\!\!- \ldots -\!\!- x_{l-1}}_{\in Q_0 \backslash A} -\!\!- \underbrace{x_l}_{\in A} -\!\!- \ldots -\!\!- \underbrace{x_{r+1} = b}_{\in Q_0 \backslash A}$$

with $j-1 \geq 0$ and $l-1 \geq j$. By the induction hypothesis there exists in $\Gamma(\mathcal{B}_Q)$ a walk between A and $A\backslash\{x_l\} \cup \{b\} =: B$. Since all $x_0, \ldots, x_{r+1}$ are pairwise disjoint the following walk is a walk with just one change (now from B to $Q_0 \backslash B$):

$$\underbrace{a -\!- x_1 -\!- \ldots -\!- x_{j-1}}_{\in B} -\!- \underbrace{x_j -\!- \ldots -\!- x_{l-1} -\!- x_l}_{\in Q_0 \backslash B}$$

Again by the induction hypothesis there is a walk in the Gabriel quiver between B and $B\backslash\{a\} \cup \{x_l\} = A\backslash\{a\} \cup \{b\}$ which finishes our walk from A to $A\backslash\{a\} \cup \{b\}$. □

Proof of Proposition 1.3. As we observed above, the Gabriel quiver of $\mathcal{B}_{Q_n}$ has at least $n+1$ connected components. Assertion (a) follows from the last lemma. Now let Q be distinct from any Q_n. Therefore Q has a subquiver of the form $x_1 \to s \leftarrow x_2$ or $x_1 \leftarrow s \to x_2$. In each case the subsets $\{x_1, x_2\}$ and $\{s\}$ are strongly connected. Consequently there is an arrow in the Gabriel quiver between $\{x_1, x_2\}$ and $\{s\}$. Thus $\Gamma(\mathcal{B}_Q)_2$ and $\Gamma(\mathcal{B}_Q)_1$ are connected. The involution ι yields an arrow in $\Gamma(\mathcal{B}_Q)$ between the subsets $Q_0\backslash\{x_1, x_2\}$ and $Q_0\backslash\{s\}$, which connects $\Gamma(\mathcal{B}_Q)_{n-2}$ and $\Gamma(\mathcal{B}_Q)_{n-1}$. Now for each subset $D \subseteq Q_0\backslash\{x_1, x_2, s\}$ there is an arrow in $\Gamma(\mathcal{B}_Q)$ between $D \cup \{x_1, x_2\}$ and $D \cup \{s\}$ linking the subquivers $\Gamma(\mathcal{B}_Q)_{|D|+1}$ and $\Gamma(\mathcal{B}_Q)_{|D|+2}$. Hence $\Gamma(\mathcal{B}_Q)_1, \ldots, \Gamma(\mathcal{B}_Q)_{n-1}$ are connected. Meanwhile $\Gamma(\mathcal{B}_Q)_0 = \bullet_\emptyset$ and $\Gamma(\mathcal{B}_Q)_{|Q_0|} = \bullet_{Q_0}$. □

5 The monoid algebra attached to $\mathbf{Q_n}$

In this section let R be a field. Recall that $\mathcal{P}_n$ denotes the poset of the product order $\geq_n$ defined in the introduction. We first introduce the elements of $\mathcal{A}_n := R\pi_{Q_n}$ to state the precise isomorphism between the incidence algebra $\mathrm{Inc}(\mathcal{P}_n)$ and $\mathcal{A}_n$. Then we prove the stated properties in the subsections 5.1, 5.2 and 5.3.

As is well-known, the incidence algebra $\mathrm{Inc}(\mathcal{P}_n)$ is the free R-space over $X_{(J,I)}$ with $J \geq_n I$ endowed with the multiplication given by $X_{(K,J)}X_{(J',I)} = X_{(K,I)}$ if $J = J'$ and $X_{(K,J)}X_{(J',I)} = 0$ otherwise. It is the path algebra of the Hasse diagramm of the poset $\mathcal{P}_n$ modulo the ideal which is generated by the commutativity relations (e.g. see [Rin84]).

As we have already seen $\mathcal{A}_1 \subseteq \mathcal{A}_2 \subseteq \mathcal{A}_3 \subseteq \ldots$ is a tower of algebras. Thus an inductive description of elements is possible. Local properties such as idempotency or orthogonality are preserved, in contrast to global ones such as centrality or being the unit element 1 in $\mathcal{A}_n$, when viewing elements in a greater algebra $\mathcal{A}_{n+k}$.

The heart of the definition of the idempotents in $\mathcal{A}_n$ corresponding to the idempotents $X_{(J,J)}$ in $\mathrm{Inc}(\mathcal{P}_n)$ are the inductively (on n) defined elements $y_1^{(n)}, y_2^{(n)}, \ldots, y_n^{(n)}$ in $\mathcal{A}_n$. We start with $y_1^{(1)} := \mathsf{P}_1$ and set for all $k \in \underline{n}\rfloor$:

$$y_k^{(n)} := \begin{cases} y_1^{(n-1)} - \mathsf{P}_n y_1^{(n-1)} + \mathsf{P}_n & \text{if } k = 1 \\ y_k^{(n-1)} - \mathsf{P}_n y_k^{(n-1)} + y_{k-1}^{(n-1)} \mathsf{P}_n & \text{if } 2 \leq k \leq n \\ 0 & \text{if } k > n \end{cases}$$

Each of these elements generates an ideal which is closely related to the admissible normal form given in 3.7 (see Corollary 5.5). Their properties are listed in Lemma 5.4. From them we conclude that the idempotents in $\mathcal{A}_n$ corresponding to the connected components of $\mathrm{Inc}\,(\mathcal{P}_n)$ are given by:

$$z_0^{(n)} := 1 - y_1^{(n)}, \quad \dots, \quad z_k^{(n)} := y_k^{(n)} - y_{k+1}^{(n)}, \quad \dots, \quad z_n^{(n)} := y_n^{(n)} - y_{n+1}^{(n)} = y_n^{(n)}$$

Furthermore we consider the inductively defined elements $g_J^{(n)} \in \mathcal{A}_n$ for all $\emptyset \neq J \subseteq \underline{n\rfloor}$ starting with $g_{\{1\}}^{(1)} := \mathsf{P}_1$:

$$g_J^{(n)} := \begin{cases} g_J^{(n-1)} - \mathsf{P}_n g_J^{(n-1)} & \text{if } n \notin J \\ \mathsf{P}_n & \text{if } \{n\} = J \\ g_{J\setminus\{n\}}^{(n-1)} \mathsf{P}_n & \text{if } n \in J \neq \{n\} \end{cases}$$

Theorem 5.1. *A complete system of pairwise orthogonal idempotents of $\mathcal{A}_n$ consists of*

$$f_\emptyset^{(n)} := z_0^{(n)} \quad \textit{and} \quad f_J^{(n)} := g_J^{(n)} z_{|J|}^{(n)} \quad \textit{with} \quad \emptyset \neq J \subseteq \underline{n\rfloor}$$

(In [DHST11] a complete set of orthogonal idempotents for $\mathcal{A}_n$ is given explicitly.) Now we turn towards those elements in $\mathcal{A}_n$ corresponding to $X_{(J,I)} \in \mathrm{Inc}\,(\mathcal{P}_n)$ with $J \neq I$. For this we define for all $n \geq j \geq i \geq 1$ the word:

$$j \leftharpoonup i := j\,.\,(j-1)\,.\,\dots\,.\,(i+1)\,.\,i \qquad \in \underline{n\rfloor}^*$$

and for all subsets $J = j_r > \dots > j_1$ and $K = k_r > \dots > k_1$ of $\underline{n\rfloor}$ with $K \geq_n J$ the monomial

$$\mathsf{P}_{(K,J)} := \mathsf{P}_{k_1 \leftharpoonup j_1 \,.\, \dots \,.\, k_r \leftharpoonup j_r} \qquad \in \mathcal{A}_n$$

The already introduced idempotent $\mathsf{P}_J = \mathsf{P}_{j_1\,.\,j_2\,.\,\dots\,.\,j_r}$ coincides with $\mathsf{P}_{(J,J)}$; meanwhile $\mathsf{P}_{(\emptyset,\emptyset)} = 1$ in $\mathcal{A}_n$. The main difficulty is to show that the elements $f_K^{(n)} \mathsf{P}_{(K,J)} f_J^{(n)}$ are distinct from 0.

Theorem 5.2. *For every pair $K \geq_n J$ we have: $f_K^{(n)} \mathsf{P}_{(K,J)} f_J^{(n)} \neq 0$.*

Here we will need two more descriptions of the middle factor, including one inductive one (see subsection 5.2.2) and an inductive description of the elements $g_J^{(n)}$ (see Lemma 5.15). Along the way we get all we need to prove the main theorem:

Main Theorem 5.3. *Let R be a field and $n \in \mathbb{N}$. The R-linear map $\Phi\colon \mathrm{Inc}\,(\mathcal{P}_n) \longrightarrow \mathcal{A}_n$ with $\Phi(X_{(K,J)}) = f_K^{(n)} \mathsf{P}_{(K,J)} f_J^{(n)}$ for all $K \geq_n J$ is an isomorphism of algebras.*

One remark on the field R: in our proof we distinguish between two monomials P_v and P_w by comparing their action on the injective indecomposable Q_n representations $I_0, I_1 \dots I_n$ over R. These steps can be replaced by comparing the non-decreasing parking functions π_v and π_w as functions on $1, \dots, n+1$ (notation as in [HT09]). Since we just use the defining relations for π_{Q_n} (see Proposition 3.7) which are the same as for NDPF_{n+1} (see [HT09]), we could replace the field R by an arbitrary commutative ring, as mentioned in [HT09]) and [DHST11]:

Remark 1. *Let R be a commutative ring. Then the R-algebras $R\mathrm{NDPF}_{n+1}$ and $\mathrm{Inc}\,(\mathcal{P}_n)$ are isomorphic.*

5.1 Proof of Theorem 5.1

We start with the properties of the elements $y_1^{(n)}, \dots, y_n^{(n)}$ and the direct conclusions for the elements $z_0^{(n)}, \dots, z_n^{(n)}$:

Lemma 5.4. *The elements* $y_j^{(n)}$ *are central in* $\mathcal{A}_n$ *for all* $j \in \underline{n}\rfloor$. *Thus the elements* $z_j^{(n)}$ *are central. Meanwhile the idempotency and orthogonality of* $z_0^{(n)}, \dots, z_n^{(n)}$ *follows directly from:*

$$y_i^{(n)} y_j^{(n)} = y_j^{(n)} \qquad \textit{for all} \quad 0 < i \leq j \leq n$$

Each $y_j^{(n)}$ *is distinct from* 0 *since for all subsets* $J \subseteq \underline{n}\rfloor$ *with* $|J| = j$ *the following equation holds:*

$$y_j^{(n)} \mathsf{P}_J = \mathsf{P}_J$$

Proof. Exemplary in more detail, we prove by induction on n that $y_i^{(n)}$ is central for each $i \in \underline{n}\rfloor$. For $n \in \{1, 2\}$ the centrality of $y_1^{(1)} = \mathsf{P}_1$ and of $y_1^{(2)} = \mathsf{P}_1 + \mathsf{P}_2 - \mathsf{P}_{2.1}$ or $y_2^{(2)} = \mathsf{P}_{1.2}$ in $\mathcal{A}_1$ resp. in $\mathcal{A}_2$ are direct consequences of the defining relations. So let $n > 2$. For a generator P_j of $\mathcal{A}_n$ we have:

$$\mathsf{P}_j y_1^{(n)} - y_1^{(n)} \mathsf{P}_j = (\mathsf{P}_j y_1^{(n-1)} - y_1^{(n-1)} \mathsf{P}_j) + (-\mathsf{P}_j \mathsf{P}_n y_1^{(n-1)} + \mathsf{P}_n y_1^{(n-1)} \mathsf{P}_j) + (\mathsf{P}_j \mathsf{P}_n - \mathsf{P}_n \mathsf{P}_j)$$

For $j < n - 1$ this adds up to 0 by the induction hypothesis and the (commutativity) relations. We consider the cases $j \in \{n - 1, n\}$ separately using the induction hypothesis for $j = n - 1$:

$$\begin{aligned}
\mathsf{P}_{n-1} y_1^{(n)} - y_1^{(n)} \mathsf{P}_{n-1} &= (-\mathsf{P}_{n-1}\mathsf{P}_n + \mathsf{P}_n \mathsf{P}_{n-1}) y_1^{(n-1)} + (\mathsf{P}_{n-1}\mathsf{P}_n - \mathsf{P}_n \mathsf{P}_{n-1}) \\
&= (-\mathsf{P}_{n-1}\mathsf{P}_n + \mathsf{P}_n \mathsf{P}_{n-1}) y_1^{(n-2)} - (-\mathsf{P}_{n-1}\mathsf{P}_n + \mathsf{P}_n \mathsf{P}_{n-1}) \mathsf{P}_{n-1} y_1^{(n-2)} \\
&\quad + (-\mathsf{P}_{n-1}\mathsf{P}_n + \mathsf{P}_n \mathsf{P}_{n-1}) \mathsf{P}_{n-1} + (\mathsf{P}_{n-1}\mathsf{P}_n - \mathsf{P}_n \mathsf{P}_{n-1}) \\
&= (-\mathsf{P}_{n-1}\mathsf{P}_n + \mathsf{P}_n \mathsf{P}_{n-1}) y_1^{(n-2)} - (-\mathsf{P}_{n-1}\mathsf{P}_n + \mathsf{P}_n \mathsf{P}_{n-1}) y_1^{(n-2)} \\
&= 0
\end{aligned}$$

and the generalized relations for $j = n$:

$$\mathsf{P}_n y_1^{(n)} - y_1^{(n)} \mathsf{P}_n = (\mathsf{P}_n y_1^{(n-1)} - y_1^{(n-1)} \mathsf{P}_n) + (-\mathsf{P}_n y_1^{(n-1)} + y_1^{(n-1)} \mathsf{P}_n) = 0$$

Now we consider the elements $y_2^{(n)}, \dots, y_n^{(n)}$. For any $k > 1$ and a generator P_j of $\mathcal{A}_n$ we have:

$$\mathsf{P}_j y_k^{(n)} - y_k^{(n)} \mathsf{P}_j = (\mathsf{P}_j y_k^{(n-1)} - y_k^{(n-1)} \mathsf{P}_j) + (-\mathsf{P}_j \mathsf{P}_n y_k^{(n-1)} + \mathsf{P}_n y_k^{(n-1)} \mathsf{P}_j) + (\mathsf{P}_j y_{k-1}^{(n-1)} \mathsf{P}_n - y_{k-1}^{(n-1)} \mathsf{P}_n \mathsf{P}_j)$$

Again for $j < n - 1$, this adds up to 0. Meanwhile we use the induction hypothesis and the generalized relations for $j = n - 1$:

$$\begin{aligned}
\mathsf{P}_{n-1}y_k^{(n)} - y_k^{(n)}\mathsf{P}_{n-1} &= (-\mathsf{P}_{n-1}\mathsf{P}_n + \mathsf{P}_n\mathsf{P}_{n-1})(y_k^{(n-2)} - \mathsf{P}_{n-1}y_k^{(n-2)} + y_{k-1}^{(n-2)}\mathsf{P}_{n-1}) \\
&\quad + (y_{k-1}^{(n-2)} - \mathsf{P}_{n-1}y_{k-1}^{(n-2)} + y_{k-2}^{(n-2)}\mathsf{P}_{n-1})(\mathsf{P}_{n-1}\mathsf{P}_n - \mathsf{P}_n\mathsf{P}_{n-1}) \\
&= (-\mathsf{P}_{n-1}\mathsf{P}_n + \mathsf{P}_n\mathsf{P}_{n-1})y_{k-1}^{(n-2)}\mathsf{P}_{n-1} + (y_{k-1}^{(n-2)} - \mathsf{P}_{n-1}y_{k-1}^{(n-2)})(\mathsf{P}_{n-1}\mathsf{P}_n - \mathsf{P}_n\mathsf{P}_{n-1}) \\
&= -\mathsf{P}_{n-1}\mathsf{P}_n y_{k-1}^{(n-2)}\mathsf{P}_{n-1} + \mathsf{P}_n y_{k-1}^{(n-2)}\mathsf{P}_{n-1} + y_{k-1}^{(n-2)}\mathsf{P}_{n-1}\mathsf{P}_n - \mathsf{P}_n y_{k-1}^{(n-2)}\mathsf{P}_{n-1} \\
&\quad - y_{k-1}^{(n-2)}\mathsf{P}_{n-1}\mathsf{P}_n + \mathsf{P}_{n-1}\mathsf{P}_n y_{k-1}^{(n-2)}\mathsf{P}_{n-1} \\
&= 0
\end{aligned}$$

For $j = n$ the calculation is again simply:

$$\mathsf{P}_n y_k^{(n)} - y_k^{(n)}\mathsf{P}_n = (\mathsf{P}_n y_k^{(n-1)} - y_k^{(n-1)}\mathsf{P}_n) + (-\mathsf{P}_n y_k^{(n-1)} + y_k^{(n-1)}\mathsf{P}_n) + (y_{k-1}^{(n-1)}\mathsf{P}_n - y_{k-1}^{(n-1)}\mathsf{P}_n) = 0$$

Now we prove by induction that $y_j^{(n)}y_k^{(n)} = y_k^{(n)}$ holds for all $1 \leq j \leq k \leq n$. The case for $n = 1$ is trivial, so we proceed with $n > 1$. For this we set $y_0^{(n-1)} := 1 \in \mathcal{A}_n$ and rewrite:

$$y_j^{(n)} = y_j^{(n)} - \mathsf{P}_n y_j^{(n-1)} + y_{j-1}^{(n-1)}\mathsf{P}_n = (1 - \mathsf{P}_n)y_j^{(n)} + y_{j-1}^{(n-1)}\mathsf{P}_n \in \mathcal{A}_n$$

By the generalized relations we get the following equalities in $\mathcal{A}_n$; the last one by the induction hypothesis:

$$\begin{aligned}
y_j^{(n)}y_k^{(n)} &= (1 - \mathsf{P}_n)y_j^{(n-1)}(1 - \mathsf{P}_n)y_k^{(n-1)} + (1 - \mathsf{P}_n)y_j^{(n-1)}y_{k-1}^{(n-1)}\mathsf{P}_n \\
&\quad + y_{j-1}^{(n-1)}\mathsf{P}_n(1 - \mathsf{P}_n)y_k^{(n-1)} + y_{j-1}^{(n-1)}\mathsf{P}_n y_{k-1}^{(n-1)}\mathsf{P}_n \\
&= (1 - \mathsf{P}_n)y_j^{(n-1)}(1 - \mathsf{P}_n)y_k^{(n-1)} + y_{j-1}^{(n-1)}y_{k-1}^{(n-1)}\mathsf{P}_n \\
&= (1 - \mathsf{P}_n)y_k^{(n-1)} + y_{k-1}^{(n-1)}\mathsf{P}_n
\end{aligned}$$

For the last statement it is convenient to show by induction on n (simultanously) that for all $\emptyset \neq J \subseteq \underline{n}\rfloor$ the following two equations hold:

$$y_{|J|}^{(n)}\mathsf{P}_J = \mathsf{P}_J \qquad \text{and} \qquad y_{|J|}^{(n)}\mathsf{P}_{n+1}\mathsf{P}_J = \mathsf{P}_{n+1}\mathsf{P}_J$$

□

Proof of Theorem 5.1. By similar computations and case by case analysis we deduce the Theorem with the following steps: first show by induction on n, that for each $k \in \underline{n}\rfloor$ the set $\{g_J^{(n)} \mid J \subseteq \underline{n}\rfloor, |J| = k\}$ is a set of pairwise orthogonal idempotents, whose elements add up to $y_k^{(n)}$. Therefore – by definition of $f_J^{(n)}$ and the properties of $z_k^{(n)}$ – the set $\{f_J^{(n)} \mid J \subseteq \underline{n}\rfloor, |J| = k\}$ consists of pairwise orthogonal idempotents. From $y_k^{(n)}z_k^{(n)} = z_k^{(n)}$ we get $\sum_{J \subseteq \underline{n}\rfloor, |J|=k} f_J^{(n)} = z_k^{(n)}$. Since $\{z_k^{(n)} \mid 0 \leq k \leq n\}$ consists of central pairwise orthogonal elements, which add up to $1 \in \mathcal{A}_n$, Theorem 5.1 follows. □

We finish this subsection with a useful remark on the following chain of ideals in $\mathcal{A}_n$:

$$\mathcal{A}_n = \mathfrak{I}_0^{(n)} \supset \mathfrak{I}_1^{(n)} \supset \mathfrak{I}_2^{(n)} \supset \ldots \supset \mathfrak{I}_{n-1}^{(n)} \supset \mathfrak{I}_n^{(n)} = \langle \mathsf{P}_{\underline{n}\rfloor} \rangle_R \supset 0$$

where $\Im_k^{(n)}$ is the ideal generated by the monomials P_J with $J \subseteq \underline{n}\rfloor$ and $|J| = k$. It is directly deduced from the last equation of Lemma 5.4, that $\Im_k^{(n)}$ is contained in the ideal $y_k^{(n)}\mathcal{A}_n$. On the other hand an easy induction shows that $y_k^{(n)}$ is contained in $\Im_k^{(n)}$. Thus the equalities hold:

Corollary 5.5.

$$\Im_k^{(n)} = y_k^{(n)}\mathcal{A}_n = \{a \in \mathcal{A}_n \mid y_k^{(n)}a = a\}$$

In particular $z_k^{(n)} = y_k^{(n)} - y_{k+1}^{(n)}$ *annihilates the ideals* $\Im_{k+1}^{(n)} \supset \ldots \supset \Im_n^{(n)}$.

5.2 Proof of Theorem 5.2

5.2.1 Chain description of $\mathsf{P}_{\mathbf{K},\mathbf{J}}$

For $n \in \mathbb{N}$ and subsets J and K of $\underline{n}\rfloor$ we write $K \gtrdot J$ if K is a minimal proper successor of J w.r.t. $\geq_n$ and say that their are neighbours. In general there is more than one $\gtrdot$-chain between $K \geq_n J$. But the monomial $\mathsf{P}_{(K,J)}$ is an invariant of such $\gtrdot$-chains:

Lemma 5.6. *Let* $K \geq_n J$. *For every* $\gtrdot$*-chain* $K = H_t \gtrdot \ldots \gtrdot H_1 = J$ *we have:*

$$\mathsf{P}_{(K,J)} = \mathsf{P}_{H_t \,.\ldots.\, H_1}$$

Proof. We just show the induction step $2 \leq n \to n+1$ for non-empty, proper subsets $n+1 \in K \geq_{n+1} J$ of $\underline{n+1}\rfloor$. Let $1 < r := |J| \neq J$ and consider a $\gtrdot$-chain $K = H_t \gtrdot \ldots \gtrdot H_1 = J$. Let $h_{r,s} > h_{r-1,s} > \ldots > h_{1,s}$ be the elements of H_s for $s \in \underline{t}\rfloor$. Since $H_{s+1} \gtrdot H_s$ there is exactly one index $k \in \underline{r}\rfloor$ with $h_{k,s+1} > h_{k,s} = h_{k,s+1} - 1$ and $h_{j,s} = h_{j,s+1}$ for all $j \neq k$.
We denote by $\widetilde{J}$, $\widetilde{K}$ and $\widetilde{H_s}$ the sets J, K and H_s without their maximal elements respectively.
First case: $n+1 \in J$ In particular $k_r \leftharpoonup j_r = n+1$ holds. Moreover the maximal elements $h_{r,s} = n+1$ for $s \in \underline{t}\rfloor$ are not involved in the $\gtrdot$-chain, that is we already have a $\gtrdot$-chain:

$$\widetilde{K} = K\backslash\{n+1\} = \widetilde{H_t} \gtrdot \ldots \gtrdot \widetilde{H_1} = J\backslash\{n+1\} = \widetilde{J}$$

So by the generalized relations and the induction hypothesis we deduce:

$$\mathsf{P}_{H_t \,.\ldots.\, H_1} = \mathsf{P}_{\widetilde{H_t}\,.\,n+1 \,.\ldots.\, \widetilde{H_2}\,.\,n+1\,.\,\widetilde{H_1}\,.\,n+1} = \mathsf{P}_{\widetilde{H_t} \,.\ldots.\, \widetilde{H_2}\,.\,\widetilde{H_1}\,.\,n+1} = \mathsf{P}_{k_1 \leftharpoonup j_1 \,.\ldots.\, k_{r-1} \leftharpoonup j_{r-1}\,.\,k_r \leftharpoonup j_r} = \mathsf{P}_{(K,J)}$$

Second case: $n+1 \notin J$ We divide the chain into two chains, such that one of them contains no $n+1$ but the other does. More precisely, there is an index $s \in \{2, \ldots, t\}$ minimal with $n+1 \in H_s$. Then we have: $h_{r,s-1} = n < n+1 = h_{r,s} = \ldots = h_{r,t} = k_r$. As in the first case we receive:

$$\mathsf{P}_{H_t \ldots H_s} = \mathsf{P}_{(\widetilde{H_s},\widetilde{H_t})}\mathsf{P}_{n+1} \quad \text{and} \quad \mathsf{P}_{H_{s-1}\ldots H_1} = \mathsf{P}_{(\widetilde{H_{s-1}},\widetilde{H_1})}\mathsf{P}_{h_{r,s-1} \leftharpoonup h_{r,1}}$$

Now P_{n+1} commutes with all P_j for $j \leq h_{r-1,s-1}$, hence with $\mathsf{P}_{(\widetilde{H_{s-1}},\widetilde{H_1})}$. Moreover we observe $\widetilde{H}_{s-1} = \widetilde{H}_s$. Therefore we get by applying the induction hypothesis several times:

$$\mathsf{P}_{H_t\ldots H_1} = \mathsf{P}_{(\widetilde{H_t},\widetilde{H_s})}\mathsf{P}_{(\widetilde{H_{s-1}},\widetilde{H_1})}\mathsf{P}_{n+1\,.\,h_{r,s-1} \leftharpoonup h_{r,1}} = \mathsf{P}_{(\widetilde{H_t},\widetilde{H_1})}\mathsf{P}_{n+1 \leftharpoonup j_r} = \mathsf{P}_{(K,J)}$$

□

Corollary 5.7. *For all* $L \geq_n K \geq_n J$ *the monomial* $\mathsf{P}_{(K,J)}$ *is contained in the ideal* $\Im_{|J|}^{(n)}$ *and we have* $\mathsf{P}_{(L,K)}\mathsf{P}_{(K,J)} = \mathsf{P}_{(L,J)}$, *in particular* $\mathsf{P}_K\mathsf{P}_{(K,J)} = \mathsf{P}_{(K,J)} = \mathsf{P}_{(K,J)}\mathsf{P}_J$ *holds.*

In fact, the stronger assertion holds:

Lemma 5.8.

$$\mathsf{P}_{(K,J)} \in \mathfrak{I}^{(n)}_{|J|} \setminus \mathfrak{I}^{(n)}_{|J|+1}$$

Proof. The monomials $\mathsf{P}_{(K,J)}$ for maximal K and minimal J (w.r.t. $\leq_n$) are easier to handle, i.e. $K = \{n-r+1, \ldots, n\}$ and $J = \underline{r}\rfloor$ for some $r \in \underline{n}\rfloor \cup \{0\}$. We set $\mathsf{m}^{(n)}_{[0]} := 1 \in \mathcal{A}_n$ and for each $r \in \underline{n}\rfloor$

$$\mathsf{m}^{(n)}_{[r]} := \mathsf{P}_{(\{n-r+1,\ldots,n\},\underline{r}\rfloor)} = \mathsf{P}_{n-r+1 \leftharpoonup 1 \,.\ldots.\, n-1 \leftharpoonup r-1 \,.\, n \leftharpoonup r} = \mathsf{m}^{(n-1)}_{[r-1]}\mathsf{P}_{n \leftharpoonup r}$$

Now we prove $\mathsf{m}^{(n)}_{[r]} \notin \mathfrak{I}^{(n)}_{r+1}$ by induction on n, which is by Corollary 5.5 equivalent to:

$$(1 - y^{(n)}_{r+1})\mathsf{m}^{(n)}_{[r]} \neq 0$$

In the induction step $n \to n+1$ the extreme cases $r \in \{0, n+1\}$ are trivial and for the remaining r we get by direct calculations using the generalized relations:

$$(1 - y^{(n+1)}_2)\mathsf{m}^{(n+1)}_{[1]} = \mathsf{z}^{(n)}_0 \mathsf{P}_{n+1 \leftharpoonup 1} \quad \text{and} \quad (1 - y^{(n+1)}_{r+1})\mathsf{m}^{(n+1)}_{[r]} = (1 - y^{(n)}_r)\mathsf{m}^{(n)}_{[r-1]}\mathsf{P}_{n+1 \leftharpoonup r}$$

The induction hypothesis applies to the ladder cases. Thus the linear combinations $z^{(n)}_0$ and $(1 - y^{(n)}_r)\mathsf{m}^{(n)}_{[r-1]}$ of monomials in $\mathcal{A}_n$ are distinct from 0. But for any two distinct monomials P_v and P_w in $\mathcal{A}_n$ there exists an injective indecomposable Q_n-representation I_y with $\mathsf{P}_v I_y \neq \mathsf{P}_w I_y$ by the proof of Proposition 3.7. With Lemma 3.5 we conclude that thus $\mathsf{P}_v\mathsf{P}_{n+1 \leftharpoonup r}$ and $\mathsf{P}_w\mathsf{P}_{n+1 \leftharpoonup r}$ act differently on one of the Q_{n+1}-representations I_y or I_{y+1}. Therefore the monomials $\mathsf{m}^{(n+1)}_{[r]}$ do not lie in $\mathfrak{I}^{(n+1)}_{r+1}$. Since each monomial $\mathsf{P}_{(K,J)}$ is a factor of $\mathsf{m}^{(n)}_{[|J|]}$ by the chain description, $\mathsf{P}_{(K,J)}$ does not lie in $\mathfrak{I}^{(n)}_{|J|+1}$. □

5.2.2 Inductive description of $\mathsf{P}_{(\mathbf{K},\mathbf{J})}$

We denote by $K_{\max}$ the greatest interval (w.r.t. $\succ$, see page 9) of a finite subset K of $\mathbb{N}$, e.g. $\{9,8,5,4,2\}_{\max} = \{9,8\}$ and $\{4,3,2\}_{\max} = \{4,3,2\}$ and $\{5,3,2,1\}_{\max} = \{5\}$. As usual we set $K - 1 := \{k - 1 \mid k \in K\}$ and $\emptyset - 1 := \emptyset$. If $K \geq_{n+1} J$ and $n+1$ lies in J, then $n+1$ also lies in K, moreover $J_{\max}$ is a subset of $K_{\max}$. Therefore the distinction of cases in the next remark is complete.

Remark 1. *Let $K \geq_{n+1} J$ and $K \neq J$. Then we have:*

$$\begin{array}{ll} K \geq_n J & \textit{if } n+1 \notin K \\ K \setminus K_{\max} \cup (K_{\max} - 1) \geq_n J & \textit{if } n+1 \in K \setminus J \\ K \setminus K_{\max} \cup ((K_{\max} \setminus J_{\max}) - 1) \geq_n J \setminus J_{\max} & \textit{if } n+1 \in J \end{array}$$

Thus we can define:

Definition 5.9. *Starting with* $\mathsf{m}^{(1)}_{[\{1\},\{1\}]} := \mathsf{P}_1$ *and* $\mathsf{m}^{(1)}_{[\emptyset,\emptyset]} = 1 \in \mathcal{A}_1$ *we define inductively on* n *for every pair* $K \geq_{n+1} J$ *the monomial* $\mathsf{m}^{(n+1)}_{[K,J]}$ *by:*

$$\mathsf{m}^{(n+1)}_{[K,J]} := \begin{cases} \mathsf{P}_J & \text{if } J = K \\ \mathsf{m}^{(n)}_{[K,J]} & \text{if } J \neq K, n+1 \notin K \\ \mathsf{P}_{K_{\max}}\ \mathsf{m}^{(n)}_{[K\setminus K_{\max}\cup(K_{\max}-1)\ ,\ J]} & \text{if } J \neq K, n+1 \in K\setminus J \\ \mathsf{P}_{K_{\max}}\ \mathsf{m}^{(n)}_{[K\setminus K_{\max}\cup((K_{\max}\setminus J_{\max})-1)\ ,\ J\setminus J_{\max}]} & \text{if } J \neq K, n+1 \in J \end{cases}$$

An induction on n and a case by case analysis according to the definition of $\mathsf{m}^{(n)}_{[K,J]}$ shows in a straightforward way:

Lemma 5.10. *Let* $K \geq_n J$. *Then we have:*

$$\mathsf{P}_{(K,J)} = \mathsf{m}^{(n)}_{[K,J]}$$

□

Before we use the inductive description of $\mathsf{P}_{(K,J)}$ we need to introduce two more notation. With them we can formulate an inductive description of $g^{(n)}_J$ and hence of $f^{(n)}_J$ (see Lemma 5.15).

Definition 5.11. *For a subset* N *of* $\underline{n}\rfloor$ *we define the element* $y(N)$ *in* $\mathcal{A}_n$ *by*

$$y(N) = \begin{cases} 0 & \text{if } N = \emptyset \\ y(N\setminus \max N) - \mathsf{P}_{\max N}\, y(N\setminus \max N) + \mathsf{P}_{\max N} & \text{if } N \neq \emptyset \end{cases}$$

So $y(N)$ is similarly defined to $y^{(n)}_1$ and has similar properties (w.r.t. to the subalgebra $\mathcal{A}_N$ of $\mathcal{A}_n$ generated by P_s with $s \in N$), namely: the element $y(N)$ is central in $\mathcal{A}_N$. Moreover, for all $m \in N$ and each $x > \max N$ we have the identities $y(N)\mathsf{P}_m = \mathsf{P}_m$ and $y(N)\mathsf{P}_x\mathsf{P}_m = \mathsf{P}_x\mathsf{P}_m$ and consequently $y(N) \neq 0$. We consider these elements for the sets:

Definition 5.12. *We define for each subset* K *of* $\underline{n}\rfloor$ *the subset* $N^{(n)}_K$ *of* $\underline{n}\rfloor$ *by:*

$$N^{(n)}_K := \begin{cases} \emptyset & \text{if } K = \emptyset \\ \{x \in \underline{n}\rfloor \setminus K \mid x > \min K\} & \text{if } K \neq \emptyset \end{cases}$$

Some examples are: $N^{(5)}_{\{1,2\}} = \{3,4,5\} = N^{(5)}_{\{2\}}, N^{(5)}_{\{1,3\}} = \{2,4,5\}, N^{(5)}_{1,3,4,5} = \{2\}$ and $N^{(5)}_{\{4,5\}} = \emptyset$.

Lemma 5.13. *For each subset* J *of* $\underline{n}\rfloor$ *the linear combination* $\mathsf{P}_J y(N^{(n)}_J)\mathsf{P}_J$ *of monomials in* $\mathcal{A}_n$ *lies in the ideal* $\Im^{(n)}_{|J|+1}$ *and is thus annihilated by* $z^{(n)}_{|J|}$ *(see Corollary 5.5).*

Proof. By Corollary 5.5 it suffices to show the identity:

$$y^{(n)}_{|J|+1}\mathsf{P}_J\, y(N^{(n)}_J)\mathsf{P}_J = \mathsf{P}_J\, y(N^{(n)}_J)\mathsf{P}_J$$

This is a straightforward induction on n requiring a case-by-case analysis on the cardinality of J and considering the cases $n+1 \in J$ and $n+1 \notin J$ separately. One also needs the identity $y^{(n)}_j\mathsf{P}_J = \mathsf{P}_J$ (see Lemma 5.4). □

The next rather technical lemma, which we prove in more detail, is the heart of the proof of Theorem 5.2. Recall that if W_n as in Proposition 3.7, then $\{\mathsf{P}_w \mid w \in W_n\}$ is a basis of $\mathcal{A}_n$.

Lemma 5.14. *Let $K \geq_n J$. Then we have:*

$$y(N_K^{(n)})\,\mathsf{P}_{(K,J)} \in \left\langle \mathsf{P}_w \in \mathcal{A}_n \;\middle|\; w \in W_n \text{ and } \mathsf{P}_v \neq \mathsf{m}^{(n)}_{[K,J]} \right\rangle_R =: U^{(n)}_{K,J}$$

Proof. The proof is an induction on n.
We start with a remark on the two extreme cases $N_K^{(n)} = \emptyset$ and $J = K$ with $N_K^{(n)} \neq \emptyset$: in the first case $y(N_K^{(n)}) = 0$ holds, so the statement is clear. In the second case let $y(N_K^{(n)}) = \sum_{w \in W_n} c_w \mathsf{P}_w$. Then we have $\mathsf{m}^{(n)}_{[K,K]}\, y(N_K^{(n)}) = \sum_{w \in W_n} c_w \mathsf{P}_K \mathsf{P}_w$. Now for all $w \in W_n$ with $c_w \neq 0$ (i.e. $\{w\} \subseteq N_K^{(n)} \subset \underline{n}\rfloor \setminus K$) the functors $\mathsf{m}^{(n)}_{[K,K]} = \mathsf{P}_K$ and $\mathsf{P}_K \mathsf{P}_w$ differ in their action on the injective indecomposable Q_n-representations I_i with $i \in \underline{n}\rfloor \setminus K$. Hence $\mathsf{m}^{(n)}_{[K,K]}\, y(N_K^{(n)})$ lies in $U^{(n)}_{K,K}$.
Since the calculations for $n \in \{1, 2, 3\}$ are trivial we proceed with the induction step $n \longrightarrow n+1$ for $n \geq 3$. Let $K \geq_{n+1} J$ such that $J \neq K$ and $N_K^{(n+1)} \neq \emptyset$ hold.
1st case: $n + 1 \notin K$. Since $N_K^{(n+1)} = N_K^{(n)} \cup \{n+1\}$ we have

$$y(N_K^{(n+1)})\mathsf{m}^{(n+1)}_{[K,J]} = \left(y(N_K^{(n)}) - \mathsf{P}_{n+1} y(N_K^{(n)}) + \mathsf{P}_{n+1} \right) \mathsf{m}^{(n)}_{[K,J]} = y(N_K^{(n)})\mathsf{m}^{(n)}_{[K,J]} + \mathsf{P}_{n+1}(1 - y(N_K^{(n)}))\mathsf{m}^{(n)}_{[K,J]}$$

By the induction hypothesis the first summand $y(N_K^{(n)})\mathsf{m}^{(n)}_{[K,J]}$ lies in $U^{(n)}_{K,J} \subseteq U^{(n+1)}_{K,J}$. Meanwhile each monomial appearing in the second summand starts with P_{n+1}, hence does not coincide with $\mathsf{m}^{(n+1)}_{[K,J]} = \mathsf{m}^{(n)}_{[K,J]} \in \mathcal{A}_n$. (Compare the actions on I_{n+1}.)
2nd case: $n + 1 \in K$ Then $N := N^{(n)}_{K \setminus \{n+1\}} = N_K^{(n+1)} \neq \emptyset$ holds and we have:

$$K \neq K_{\max} \qquad \text{and} \qquad \max N = \min K_{\max} - 1 \neq 0$$

We denote by $\widetilde{N}$ the set $N \setminus \{\max N\}$. Let $\widetilde{J}$ and $\widetilde{K}$ be those subsets of $\underline{n}\rfloor$ given by the definition of $\mathsf{m}^{(n+1)}_{[K,J]}$ (depending on $n + 1 \in J$ or $n + 1 \neq J$) such that we have:

$$\mathsf{m}^{(n+1)}_{[K,J]} = \mathsf{P}_{K_{\max}} \mathsf{m}^{(n)}_{[\widetilde{K},\widetilde{J}]}$$

In the sequel we show:

$$a := \mathsf{P}_{K_{\max}} y(\widetilde{N})\mathsf{m}^{(n)}_{[\widetilde{K},\widetilde{J}]} \in U^{(n+1)}_{K,J}$$
$$b := \mathsf{P}_{\max N}\mathsf{P}_{K_{\max}} \left(1 - y(\widetilde{N})\right) \mathsf{m}^{(n)}_{[\widetilde{K},\widetilde{J}]} \in U^{(n+1)}_{K,J}$$

Then the claim follows immediately because we have

$$y(N_K^{(n+1)})\mathsf{m}^{(n+1)}_{[K,J]} = \left((1 - \mathsf{P}_{\max N}) y(\widetilde{N}) + \mathsf{P}_{\max N} \right) \mathsf{P}_{K_{\max}} \mathsf{m}^{(n)}_{[\widetilde{K},\widetilde{J}]} = a + b$$

Proof of $b \in U^{(n+1)}_{K,J}$. Let $c_w \mathsf{P}_{\max N} \mathsf{P}_{K_{\max}} \mathsf{P}_w \neq 0$ be a summand of b. Note that $\{w\}$ is a subset of

$\underline{n}\rfloor$. We look at the action on the injective indecomposable Q_{n+1}-representation I_{n+1} to distinguish between $\mathsf{P}_{\max N}\mathsf{P}_{K_{\max}}\mathsf{P}_w$ and $\mathsf{m}^{(n+1)}_{[K,J]}$:

$$\begin{aligned}\mathsf{P}_{\max N}\mathsf{P}_{K_{\max}}\mathsf{P}_w(I_{n+1}) = \mathsf{P}_{\max N}\mathsf{P}_{K_{\max}}(I_{n+1}) &= \mathsf{P}_{\max N}(I_{\min K_{\max}-1}) = I_{\min K_{\max}-2}\\ &\neq I_{\min K_{\max}-1} = \mathsf{P}_{K_{\max}}(I_{n+1}) = \mathsf{P}_{K_{\max}}\mathsf{m}^{(n)}_{[\widetilde{K},\widetilde{J}]}(I_{n+1}) = \mathsf{m}^{(n+1)}_{[K,J]}(I_{n+1})\end{aligned}$$

Proof of $a \in U^{(n+1)}_{K,J}$. For $\widetilde{N} = \emptyset$ the summand a equals 0. So we assume $\widetilde{N} \neq \emptyset$. Now we consider:

$$s := \begin{cases} n & \text{if } n+1 \notin J \\ j := \min J_{\max} - 2 & \text{if } n+1 \in J \end{cases}$$

In each case $\widetilde{J}$ and $\widetilde{K}$ both lie in $\mathcal{A}_s$ and we have

$$\widetilde{N} = N^{(s)}_{\widetilde{K}} \qquad \text{and} \qquad \mathsf{m}^{(n)}_{[\widetilde{K},\widetilde{J}]} = \mathsf{m}^{(s)}_{[\widetilde{K},\widetilde{J}]}$$

In particular it follows:

$$a = \mathsf{P}_{K_{\max}} y(\widetilde{N})\mathsf{m}^{(n)}_{[\widetilde{K},\widetilde{J}]} = \mathsf{P}_{K_{\max}} y(N^{(s)}_{\widetilde{K}})\mathsf{m}^{(s)}_{[\widetilde{K},\widetilde{J}]}$$

Now we consider an arbitrary summand $0 \neq c_w\mathsf{P}_w = c_w\mathsf{P}_{K_{\max}}\mathsf{P}_v$ of a. By the induction hypothesis P_v and $\mathsf{m}^{(s)}_{[\widetilde{K},\widetilde{J}]}$ are distinct monomials in $\mathcal{A}_s$. Recall that by the proof of Proposition 3.7 there thus exist an injective indecomposable Q_s-representation I_j with $j \in \underline{s}\rfloor$ and an index $x \in \underline{j}\rfloor \cup \{0\}$ with:

$$I_x = \mathsf{P}_v(I_j) \neq \mathsf{m}^{(s)}_{[\widetilde{K},\widetilde{J}]}(I_j)$$

Let $u \in W_s$ with $\mathsf{m}^{(s)}_{[\widetilde{K},\widetilde{J}]} = \mathsf{P}_{\widetilde{K}_{\max}}\mathsf{P}_u$ and let $y \in \underline{j}\rfloor \cup \{0\}$ be the index such that we have:

$$I_y = \mathsf{P}_u(I_j)$$

In particular we get:

$$\mathsf{m}^{(s)}_{[\widetilde{K},\widetilde{J}]}(I_j) = \mathsf{P}_{\widetilde{K}_{\max}} I_y = \begin{cases} I_y & \text{if } y \notin \widetilde{K}_{\max} \\ I_{\min \widetilde{K}_{\max}-1} & \text{if } y \in \widetilde{K}_{\max} \end{cases}$$

Since $n+1 > y \notin \widetilde{K}_{\max}$ implies $y \notin K_{\max}$ we conclude $\mathsf{m}^{(n+1)}_{[K,J]}(I_j) = \mathsf{m}^{(s)}_{[\widetilde{K},\widetilde{J}]}(I_j)$ from:

$$\mathsf{m}^{(n+1)}_{[K,J]}(I_j) = \mathsf{P}_{K_{\max}}\mathsf{P}_{\widetilde{K}_{\max}}\mathsf{P}_u(I_j) = \begin{cases} \mathsf{P}_{K_{\max}} I_y & \text{if } y \notin \widetilde{K}_{\max} \\ \mathsf{P}_{K_{\max}} I_{\min \widetilde{K}_{\max}-1} & \text{if } y \in \widetilde{K}_{\max} \end{cases}$$

Meanwhile the action of P_w on I_j is:

$$\mathsf{P}_w(I_j) = \mathsf{P}_{K_{\max}}\mathsf{P}_v(I_j) = \mathsf{P}_{K_{\max}}(I_x) = \begin{cases} I_x & \text{if } x \notin K_{\max} \\ I_{\min K_{\max}-1} & \text{if } x \in K_{\max} \end{cases}$$

We finish the proof with a case-by-case-comparison.
$x \notin K_{\max}$: Then $\mathsf{P}_w(I_j) = \mathsf{P}_v(I_j) \neq \mathsf{m}^{(s)}_{[\widetilde{K},\widetilde{J}]}(I_j) = \mathsf{m}^{(n+1)}_{[K,J]}(I_j)$.

$\underline{x \in K_{\max}:}$ If $y \in \widetilde{K}_{\max}$ then P_w and $\mathsf{m}^{(n+1)}_{[K,J]}$ act differently on I_j since $\min \widetilde{K}_{\max} < \min K_{\max}$. So we assume $y \notin \widetilde{K}_{\max}$. We have to show, that $y \neq \min K_{\max} - 1$. The only case in which $\min K_{\max} - 1$ is not contained in $\widetilde{K}_{\max}$ is $n+1 \in J$ and $J_{\max} = K_{\max}$. But in that case $y \leq j \leq s = \min J_{\max} - 2 = \min K_{\max} - 2 < \min K_{\max} - 1$ holds. □

5.2.3 An element in $\mathbf{f^{(n)}_K}\mathcal{A}_\mathbf{n}\mathbf{f^{(n)}_J}$, Proof of Theorem 5.2

There is also an alternative description of the idempotents $g^{(n)}_J$. A straightforward induction on n now shows:

Lemma 5.15. *For each non-empty subset J of $\underline{n\rfloor}$ we have:*

$$g^{(n)}_J = \mathsf{P}_J - y(N^{(n)}_J)\mathsf{P}_J$$

In particular $g^{(n)}_J$ is contained in the ideal $\mathfrak{I}^{(n)}_k$ with $k = |J|$.

Proof of Theorem 5.2. Let $K \geq_n J$ and $k := |K| = |J|$. With the previous lemma, Lemma 5.13, we gain the first reduction of the term/sum $f^{(n)}_K \mathsf{P}_{(K,J)} f^{(n)}_J$:

$$f^{(n)}_K \mathsf{P}_{(K,J)} f^{(n)}_J = z^{(n)}_k g^{(n)}_K \mathsf{P}_{(K,J)} g^{(n)}_J = z^{(n)}_k g^{(n)}_K \mathsf{P}_{(K,J)} \mathsf{P}_J \Big(\mathsf{P}_J - y(N^{(n)}_J)\mathsf{P}_J \Big) = z^{(n)}_k g^{(n)}_K \mathsf{P}_{(K,J)}$$

From the Lemma 5.4 and the chain description of $\mathsf{P}_{(K,J)}$ we conclude next:

$$f^{(n)}_K \mathsf{P}_{(K,J)} f^{(n)}_J = y^{(n)}_k g^{(n)}_K \mathsf{P}_{(K,J)} - y^{(n)}_{k+1} g^{(n)}_K \mathsf{P}_{(K,J)} \in g^{(n)}_K \mathsf{P}_{(K,J)} + \mathfrak{I}^{(n)}_{k+1}$$

We finish the proof by looking closer at $g^{(n)}_K \mathsf{P}_{(K,J)} = \mathsf{P}_{(K,J)} - y(N^{(n)}_K)\mathsf{P}_{(K,J)}$: on the one hand we have $\mathsf{P}_{(K,J)} \notin \mathfrak{I}^{(n)}_{k+1}$ as shown in Lemma 5.8. On the other hand $\mathsf{P}_{(K,J)} - y(N^{(n)}_K)\mathsf{P}_{(K,J)} \neq 0$ holds by Lemma 5.14. Thus it follows $f^{(n)}_K \mathsf{P}_{(K,J)} f^{(n)}_J \neq 0$. □

5.3 Proof of the Main Theorem 5.3

By Theorem 5.1 and 5.2 $\{ f^{(n)}_K \mathsf{P}_{(K,J)} f^{(n)}_J \mid K \geq_n J \}$ is a linearly independent set with exactly $|\mathcal{P}_n|$ elements. Thus bijectivity follows from Proposition 3.7. To see the multiplicity let $M \geq_n L$ and $K \geq_n J$. If $K \neq L$ holds, then $f^{(n)}_L$ and $f^{(n)}_K$ are orthogonal. Hence $\Phi(X_{(M,L)})\Phi(X_{(K,J)}) = 0 = \Phi(X_{(M,L)}X_{(K,J)}) = 0$. If $K = L$ we have the following identities, the fourth identity follows from Lemma 5.13 and the fifth from the chain description):

$$\begin{aligned}
\Phi(X_{(M,L)})\Phi(X_{(K,J)}) &= f^{(n)}_M \mathsf{P}_{(M,K)} f^{(n)}_K f^{(n)}_K \mathsf{P}_{(K,J)} f^{(n)}_J \\
&= f^{(n)}_M \mathsf{P}_{(M,K)} g^{(n)}_K \mathsf{P}_{(K,J)} f^{(n)}_J \\
&= f^{(n)}_M \mathsf{P}_{(M,K)} \Big(\mathsf{P}_K - y(N^{(n)}_K) \Big) \mathsf{P}_K \mathsf{P}_{(K,J)} f^{(n)}_J \\
&= f^{(n)}_M \mathsf{P}_{(M,K)} \mathsf{P}_K \mathsf{P}_K \mathsf{P}_{(K,J)} f^{(n)}_J \\
&= f^{(n)}_M \mathsf{P}_{(M,J)} f^{(n)}_J \\
&= \Phi(X_{(M,L)}X_{(K,J)})
\end{aligned}$$

□

References

[DHST11] Tom Denton, Florent Hivert, Anne Schilling, Nicolas Thiéry, *On the representation theory of finite J-trivial monoids*, Séminaire Lotharingien de Combinatoire, B64d (2011), 44 pp.

[Fay05] Matthew Fayers, *0-Hecke algebras of finite Coxeter groups*, J. Pure Appl. Algebra **199** (2005), no. 1-3, 27–41. MR 2134290 (2005m:20016)

[GL91] Werner Geigle and Helmut Lenzing, *Perpendicular categories with applications to representations and sheaves*, Journal of Algebra **144** (1991), no. 2, 273–343.

[GM11] Olexandr Ganyushkin and Volodymyr Mazorchuk, *On Kiselman quotients of 0-Hecke monoids*, Int. Electron. J. Algebra **10** (2011), 174–191. MR 2821178

[HT09] Florent Hivert and Nicolas M. Thiéry, *The Hecke group algebra of a Coxeter group and its representation theory*, J. Algebra **321** (2009), no. 8, 2230–2258. MR 2501519 (2010a:20010)

[KM09] Ganna Kudryavtseva and Volodymyr Mazorchuk, *On Kiselman's semigroup*, Yokohama Mathematical Journal Vol. 55 (2009)

[Maz10] V. Mazorchuk, *Lectures on algebraic categorification*, ArXiv e-prints (2010).

[Nor79] P. N. Norton, 0-*Hecke algebras*, J. Austral. Math. Soc. Ser. A **27** (1979), no. 3, 337–357.

[Paa11] Anna-Louise Paasch, *Monoidalgebren von Projektionsfunktoren*, urn:nbn:de:hbz:468-20111018-152557-6, http://nbn-resolving.de/urn/resolver.pl?urn=urn018-152557-6 (2011).

[Pie82] Richard S. Pierce, *Associative algebras*, Graduate Texts in Mathematics, vol. 88, Springer-Verlag, New York, 1982, Studies in the History of Modern Science, 9.

[Pin] J.-E. Pin, *Mathematical Foundations of Automata Theory*, Preprint available at www.liafa.jussieu.fr/~jep/PDF/MPRI/MPRI.pdf

[Rin84] Claus Michael Ringel, *Tame algebras and integral quadratic forms*, Lecture Notes in Mathematics, vol. 1099, Springer-Verlag, Berlin, 1984.

[Sch91] Aidan Schofield, *Semi-invariants of quivers*, J. London Math. Soc. (2) **43** (1991), no. 3, 385–395.

[Sta99] Richard P. Stanley, *Enumerative combinatorics. Vol. 2*, Cambridge Studies in Advanced Mathematics, vol. 62, Cambridge University Press, Cambridge, 1999, With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin.